

The Oklahoma Cooperative Extension Service Bringing the University to You!

The Cooperative Extension Service is the largest, most successful informal educational organization in the world. It is a nationwide system funded and guided by a partnership of federal, state, and local governments that delivers information to help people help themselves through the land-grant university system.

Extension carries out programs in the broad categories of agriculture, natural resources and environment; family and consumer sciences; 4-H and other youth; and community resource development. Extension staff members live and work among the people they serve to help stimulate and educate Americans to plan ahead and cope with their problems.

Some characteristics of the Cooperative Extension system are:

- The federal, state, and local governments cooperatively share in its financial support and program direction.
- It is administered by the land-grant university as designated by the state legislature through an Extension director.
- Extension programs are nonpolitical, objective, and research-based information.
- It provides practical, problem-oriented education

for people of all ages. It is designated to take the knowledge of the university to those persons who do not or cannot participate in the formal classroom instruction of the university.

- It utilizes research from university, government, and other sources to help people make their own decisions.
- More than a million volunteers help multiply the impact of the Extension professional staff.
- It dispenses no funds to the public.
- It is not a regulatory agency, but it does inform people of regulations and of their options in meeting them.
- Local programs are developed and carried out in full recognition of national problems and goals.
- The Extension staff educates people through personal contacts, meetings, demonstrations, and the mass media.
- Extension has the built-in flexibility to adjust its programs and subject matter to meet new needs. Activities shift from year to year as citizen groups and Extension workers close to the problems advise changes.



David Lalman

Professor, Beef Cattle

On average, nearly 40 percent of total operating costs in cow-calf enterprises are associated with nutrition because purchased and harvested hay and concentrate feeds make up the majority of that cost. Consequently, the nutritional program represents a major target to trim cost of production. However, it is widely recognized that nutritional status of the cow is closely related to reproductive performance. If too many corners are cut in the nutritional program, pregnancy and calving rate dramatically suffer.

A ranching operation can appropriately be considered a forage production and utilization enterprise. Ranchers are in the business of converting sunlight, water and carbon dioxide into a high quality human food resource; namely beef. With good management, forage is an extremely valuable renewable resource. As such, it represents the least expensive feed resource to maintain animal health and production in cow-calf and many stocker operations. A combination of excellent forage production and grazing management practices, cattle genetics that match the forage resources, and a well-timed calving season results in minimum reliance on purchased and harvested feeds. Nevertheless, specific nutrients must be supplemented at times. While special emphasis is placed on supplementing beef cows in this section, much of the discussion and data presented are relevant to all grazing cattle.

Occasionally, cow-calf producers need to feed a concentrate or harvested forage to further increase body condition of the cows or to replace pasture forage due to limited pasture forage availability. This practice, known as feeding or substitution, is in contrast to supplementation because the alternative feed or forage actually replaces consumption of the original forage resource. In general, consumption of the original forage resource declines when cattle are fed concentrate feeds at the rate of 0.5 percent of body weight (6 pounds for 1,200-pound cows) or more. Substitute feeding is more frequently used for growing cattle than it is for mature beef cows.

Identifying a Supplemental Need

The first step in implementing and maintaining an efficient supplementation program for grazing or forage fed cattle is to identify specific supplementation needs. In other words, the producer must identify specific forage nutrients not provided in adequate quantity to meet the animal's nutrient requirements. For grazing cattle, this is a difficult task because forage quality is constantly changing and so are the animal's nutrient requirements. Knowledge of these two factors is the basis for effective and efficient supplementation. Even though the task may seem difficult, years of research and current technology provides guidance in developing an effective supplement development and evaluation plan. The following steps provide

Supplementing Beef Cows

Oklahoma Cooperative Extension Fact Sheets
are also available on our website at:
facts.okstate.edu

a logical approach in identifying a supplemental need and evaluating supplement alternatives.

1. Determine the nutrient requirements for the appropriate stage of production.
2. Estimate the amount of nutrients cows will receive from forage.
3. Subtract item No. 1 from item No. 2 to determine if a nutrient deficiency or excess exists.
4. Evaluate supplement alternatives.

It should be noted that all possible combinations of the above factors are not available in the tables, simply because there are infinite possibilities. Computer software, such as OSU Cowculator, can better pinpoint an animal's nutrient requirement at a specific time and in a specific situation. These and other useful tools can be found at beefextension.com.

Average nutrient composition of various feeds and forages common to Oklahoma are presented in Table 1. Anticipating nutrients supplied by the forage base is the most difficult task in grazing cattle nutrition. The formula for nutrient intake is simple: forage intake multiplied by concentration of available nutrients in the forage.

However, many factors influence both components in this formula. Forage intake is dramatically influenced by forage quality as well as forage availability, and both of these factors can vary dramatically from year to year and month to month.

The next step is to estimate nutrient content of standing forage or hay. As mentioned earlier, these values also vary, depending on forage type, maturity and weathering. The most accurate method to determine supplemental needs for cows that will primarily receive a hay diet is to have the hay analyzed for nutrient concentration. This will cost from \$15 to \$70 per sample, but can save hundreds, even thousands of dollars in some cases. As a starting point, Table 1 includes average nutrient values for a few common feeds and forages found in the Southern Plains.

Once nutrient requirements have been established and a reasonable estimate of the nutrient contribution of the forage has been made, determining supplemental needs is simply a comparison of the two. Again, this comparison is easily and perhaps more accurately made using computer software, such as OSU Cowculator. For this example, it is assumed cows will graze winter range with little to no hay supplementation.

Average cow weight will be 1,100 pounds and average calving date is March 15. Consequently, these cows would

Oklahoma State University, in compliance with Title VI and VII of the Civil Rights Act of 1964, Executive Order 11246 as amended, and Title IX of the Education Amendments of 1972 (Higher Education Act), the Americans with Disabilities Act of 1990, and other federal and state laws and regulations, does not discriminate on the basis of race, color, national origin, genetic information, sex, age, sexual orientation, gender identity, religion, disability, or status as a veteran, in any of its policies, practices or procedures. This provision includes, but is not limited to admissions, employment, financial aid, and educational services. The Director of Equal Opportunity, 408 Whitehurst, OSU, Stillwater, OK 74078-1035; Phone 405-744-5371; email: eeo@okstate.edu has been designated to handle inquiries regarding non-discrimination policies; Director of Equal Opportunity. Any person (student, faculty, or staff) who believes that discriminatory practices have been engaged in based on gender may discuss his or her concerns and file informal or formal complaints of possible violations of Title IX with OSU's Title IX Coordinator 405-744-9154.

Issued in furtherance of Cooperative Extension work, acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture, Director of Oklahoma Cooperative Extension Service, Oklahoma State University, Stillwater, Oklahoma. This publication is printed and issued by Oklahoma State University as authorized by the Vice President for Agricultural Programs and has been prepared and distributed at a cost of 20 cents per copy. Revised 0518 GH.

Table 1. Nutrient supply compared to requirements for an 1,100-lb beef cow grazing native range during last one third of pregnancy.

	<i>Crude Protein</i> <i>lbs per day</i>	<i>TDN</i> <i>lbs per day</i>	
Required	1.8	12.1	
Supplied by forage ¹	0.88	10.78	
Supplemental need	0.92	1.32	
	Feed Protein Concentration		
	20%	25%	38%
Amount to feed, lb	4.60	3.70	2.40
Cost per ton	300.00	340.00	380.00
Cost per day	\$0.69	\$0.63	\$0.46
Protein supplied, lb	0.92	0.93	0.91
TDN supplied, lb ¹	3.45	2.78	1.80

¹ All supplements are assumed to contain 75% TDN.

Source: NRC, 2000.

be grazing low-quality winter range throughout the last one third of gestation. Table 1 shows that this 1,100-pound cow requires about 1.8 pounds of protein and 12.1 pound of Total Digestible Nutrients (TDN) per day. These cows would be expected to consume around 2 percent of their body weight, or 22 pounds of diet dry matter (1,100 pounds x 2 percent), assuming adequate supplemental protein is provided.

By using this information, supplemental needs can be calculated as shown in Table 1. Without supplementation, this group of cows would be deficient in both protein and energy and would be expected to lose considerable body condition before calving. Here, beef cows are used in the example. However, the process to determine supplemental needs for growing cattle is the same.

Once the supplemental need is determined, various supplement alternatives are relatively easy to compare. In this example, all three supplement alternatives provide adequate protein when fed at the daily amount shown. Energy or TDN is provided in considerable excess (compared to the supplemental need) with the 20 percent supplement option. Therefore, this strategy might be desirable if increased weight gain or body condition were desired. However, if the cows were in good body condition, this strategy would simply be more expensive than one of the other strategies given in the example because of the increased feeding rate. Feeding 2.4 pounds of 38 percent supplement provides adequate protein and energy to maintain bodyweight and condition during this stage of production.

Producers can make these calculations using this approach or a computer software program, then evaluate the costs, necessary feeding rate, convenience and expected animal performance outcome for each possible alternative.

In cases where one supplemental nutrient is needed, a very effective method to evaluate cost of nutrient sources is on a cost per unit of nutrient basis. In the example, the primary nutrient needed is protein. Assuming the 20 percent supplement cost \$300 per ton, the cost per pound of protein is \$.75 (\$300 per ton divided by 400 lbs of protein per ton). If the 38 percent supplement costs \$380 per ton, the cost per pound of protein is \$.50 (\$380 per ton divided by 760 pounds of protein per ton).

In this example, the cows can maintain their current weight and body condition (assuming that it is adequate) with a supplementation program that costs about \$.46 per head per day. Had the producer chosen the 20 percent supplement program, not recognizing that the higher feeding rate and therefore higher energy intake was not necessary, the producer would spend about \$.69 per head per day or approximately \$20.70 more per cow in a 90-day period.

Additional Considerations for Supplementing Low Quality Forage

Supplementation Priorities

If supplementation is the goal for cattle grazing low quality forage, priority should first be placed on meeting the protein requirement to maximize forage intake and digestion. Many years of research have consistently shown that protein supplementation is extremely effective for cattle grazing protein-deficient forage (Table 21.2). In fact, energy supplementation will not be effective if dietary protein is deficient.

Once the producer ensures that the supplementation (or feeding) program will meet the protein requirement, energy intake should be evaluated, similar to the example given in

Table 2. Influence of winter protein supplementation on performance of beef cows grazing native range.

<i>Item</i>	<i>Treatment</i>	
	<i>2 lbs/day of 40% protein supplement</i>	<i>No supplement</i>
Cow weight change		
during late gestation	23	-153
Cow BCS ^a change		
during late gestation	-.33	-1.61
Calf birth weight	88.5	77.5
Calf weaning weight	484	448

^a Body Condition Score
Source: Steele.

References

- Aljoe, H. (2000) Limit Winter Grazing as a Protein Supplement. Ag News and Views, Noble Foundation. Retrieved from <http://www.noble.org/Ag/Forage/WinterGrazing/Index.htm>
- Altom, W., and T.F. Schmedt (1983) Limit Grazing of Small Grain Pastures. Proceedings: National Wheat Pasture Symposium. Oklahoma Agricultural Experiment Station MP-115: 297-315.
- Gill, D.R. et al. (1989) Feeding the Cow Herd. Oklahoma Beef Cattle Manual, (3rd Edition) Oklahoma Agricultural Experiment Station E-913:1-8.
- Hibberd, C.A. and T.A. Thrift (1992) Supplementation of Forage-Based Diets: Are Results Predictable? *Journal of Animal Science* 70 (Suppl. 1):181. (Abstr.).

- National Research Council. (2000) *Nutrient Requirements of Beef Cattle* (7th Edition). National Academy Press, Washington, D.C.
- Pope, L.S., Nelson, A.B., and Campbell, W.D. (1963) Feeding protein supplements to range beef cows at 2, 4, or 6-day intervals. Feeder's Day Report. Oklahoma Agricultural Experiment Station Pg. 49:51.
- Steele, J.D. et al. (2002) Drought Stressed Soybean Supplementation for Gestating Beef Cows Grazing Native Tall Grass Prairie. Oklahoma Agricultural Experiment Station Research Report P-993. Retrieved February 4, 2004 from <http://www.ansi.okstate.edu/research/2002rr/index.htm>
- Wettemann, R.P., and K.S. Lusby (1994) Influence of Interval of Feeding Protein Supplement to Spring Calving Beef Cows on Body Weight and Body Condition Score During the Winter. Oklahoma Agricultural Experiment Station Research Report P-939:123-125.

by the kidneys through urine. However, the animal is able to eliminate excess salt only when adequate clean water is available. Therefore, an abundant, clean water supply is a must when this method is used.

Salt toxicities are most likely to occur:

1. Where cattle have been deprived of salt for extended periods of time and suddenly have readily available salt.
2. Cattle eat excessive salt with an inadequate water supply.
3. When cattle drink water containing a high concentration of salt.

As a rule of thumb, cattle on salt mixtures drink 50 percent to 75 percent more water than normal, or approximately 5 gallons of additional water for each pound of salt. If only salty water is available, cattle will often refuse the supplement or may be forced into a toxicity situation. Salt content of water is usually measured by total dissolved solids (TDS) which includes calcium, magnesium, sodium chlorides, sulfates and bicarbonates. In general, caution is necessary in using salt-limited supplements when water contains more than 5,000 parts per million TDS. This analysis can usually be obtained through the Soil, Water and Forage Analysis Laboratory (SWAFL) of OSU (check with your local county Extension educator).

Salt used in self-fed supplements should be coarse, plain white salt. Cost alone prohibits the use of trace-mineralized salt; however, this should be avoided since forced feeding high levels of trace-mineralized salt could result in toxicity or mineral imbalances due to excessive intake of certain trace elements. If cattle need trace-mineralized salt, the amount consumed daily should not exceed 0.02 percent of the animal's body weight.

Controlled experiments in several states have failed to show any harmful effects upon cattle production from proper use of salt-concentrate mixes. High salt intake with adequate water has had no effect on fertility, calf crop percentage, weaning weight or appearance of animals.

Adjusting Salt Levels

Several factors influence the concentration of salt required in a mix to achieve a certain feed intake. Where large amounts of salt are naturally present in drinking water or forage, the amount of salt in the mix must be reduced to get satisfactory feed intake. On the other hand, it usually is necessary to increase the salt content of the mix through a period of time, as cattle become accustomed to the high salt level. Cattle also tend to consume more of a salt-limited supplement when forage is scarce or unpalatable. Extra precautions should be taken with these and other emergency conditions to ensure water supplies are adequate.

Estimates of salt needed to limit feed intake are shown in Table 8. Actual salt intake occasionally varies from the indicated values. Forage intake, palatability of supplement ingredients, salt content of the water and animal adaptation influence salt intake.

When cattle are accustomed to eating supplements but unaccustomed to self-feeding, prevent overeating by starting with a high salt level (50-50 or even 60-40 ratio of salt to meal).

Table 8. Estimated salt intake of cattle fed salt limited supplements^a.

Body weight, lbs	Salt consumption, lbs/day		
	Low	Average	High
300	0.3	0.5	0.6
500	0.5	0.6	0.7
700	0.6	0.7	0.9
900	0.7	0.9	1.1
1,100	0.8	1.1	1.3
1,300	0.9	1.3	1.5

^a Assumes drinking water is low in total dissolved solids (TDS).

Then, the salt level should be reduced to obtain the desired level of intake. If cattle have not eaten concentrates before, a training period of a week or more of daily hand feeding of meal without added salt may be necessary.

If grain is included in a self-fed supplement, it should be cracked or coarsely ground and mixed with salt of similar particle size. This prevents separation of the salt from the grain and aids in preventing overeating. Adequate grass or hay must be available so the cattle are not forced to eat a salt-limited supplement to survive.

Example: A producer wants to self-feed cottonseed meal at the rate of 2 pounds per head per day to a group of 1,100-pound cows. Table 8 indicates that the daily salt consumption of 1,100-pound cattle averages 1.1 pounds when salt is used to limit supplement intake. Therefore, the producer's feed blend should include 1.1 parts salt and 2 parts cottonseed meal. Total intake would be approximately 3.1 pounds per day and the blend would contain 35 percent salt. The producer will need to monitor intake and adjust these percentages slightly to achieve the desired feed intake.

Assume that in addition to 2 pounds protein supplement, it is desired that the cow also consumes 3 pounds of grain (corn, milo, etc.) for a total nonsalt consumption of 5 pounds. In this case, the blend would contain 1.1 parts salt, 2 parts cottonseed meal and 3 parts corn grain for a total of 6.1 pounds intake per day. This blend would contain 18 percent salt.

Conclusion

Reducing feed costs, while maintaining performance is a must for Oklahoma cow-calf producers. By using a systematic approach to evaluating beef cow nutritional requirements, forage nutrient contribution and alternative supplemental sources, an optimal winter nutrition program can be designed. The lowest cost alternative will not always be the best program, due to the relative value of convenience, labor availability, and feeding system. The most effective way to evaluate alternatives is to first determine the cost of the total supplementation program, then compare differences in cost with other factors.

Table 1. The decision must be made whether the cattle need to maintain body weight and condition, gain weight and condition or can be allowed to lose some weight and condition. This decision will dictate how much supplemental energy should be provided.

Lastly, vitamin and mineral requirements should be compared to expected intake, have potential deficiencies identified, and have supplemental alternatives evaluated. This is not to say that vitamins and minerals are not important. Priority is given to protein and energy nutrition first because these items are needed in much greater quantities and they have the potential to have much greater impact on animal performance and efficiency of forage utilization.

Protein Sources

Protein from plant origin (such as soybean meal, cottonseed meal, corn gluten feed, wheat middlings or alfalfa hay) generally results in better utilization of low quality roughages compared to nonprotein nitrogen sources such as urea and biuret. This is particularly true when a small amount of supplement is fed (0.5 percent of body weight or less). Nonprotein nitrogen sources are more effective in stimulating diet utilization and animal performance under one or more of the following conditions:

- When greater than 0.5 percent of body weight concentrate is being fed.
- When larger, more mature animals are being supplemented (greater than 600 pounds).
- When the protein deficiency in the diet is marginal (1 percent to 3 percent more protein needed in diet compared to 4 percent to 8 percent needed).
- When a blend of plant protein and nonprotein nitrogen sources are used.
- When it is provided in a form for animals to access more than one time per day.

Generally, when three or more of these conditions exist, studies have shown that nonprotein nitrogen sources are from 75 percent to 95 percent as effective compared to an all-natural plant protein source.

Alfalfa hay and alfalfa pellets are excellent supplements for moderate to low quality roughage growing programs. Alfalfa has long been known to have very favorable effects on rumen fermentation, and is so common in most regions of Oklahoma that it is often overlooked as an ingredient or stand-alone supplement. Studies at Kansas State University show alfalfa is equal to mixtures of grain and soybean meal containing the same percent of protein when used to supplement roughages.

Interval Feeding

Significant costs in wintering cows and stockers on dry grass are the labor and transportation required to feed supplements. Adequate research has shown cows do not need to receive protein supplements every day. In one experiment using cottonseed meal as the protein source, cows were fed the same weekly amount of supplement on two-, four-, and six-day intervals (Table 3). Although cow weight loss was slightly less when cows were fed on four-day intervals, there was no difference in cow weight loss between two- and six-day intervals. Calf weaning weights were similar among all treatments. In a more recent study, cows were fed the same amount of cottonseed meal-based protein supplement

Table 3. Performance of beef cows fed supplement at different time intervals.

	Interval between feeding, days		
	2	4	6
Supplement, lb/feeding (41% cottonseed meal)	5	10	15
Cow weight change, lb	-185	-148	-170
Calf weaning weight, lb	433	440	428

Source: Pope.

weekly, although the feeding intervals were three times per week or six times per week (Table 4). In this study, there was no difference in cow weight loss, body condition score, or pregnancy rate due to supplement feeding interval. Many ranchers follow the practice of feeding two times the daily allowance on alternate days or feeding three times per week to eliminate Sunday feeding. With interval feeding, timid cows are more likely to receive their share of supplement. Even if cows are not fed daily, they should be observed as often as necessary, especially during the calving season.

It should be noted these results were obtained using dry supplements formulated with oilseed meals. These supplements had a high concentration of plant-based protein, which has a slower rate of degradation compared to supplements containing significant amounts of nonprotein nitrogen. Cows would not be expected to perform as well if dry supplements containing significant amounts of nonprotein nitrogen were fed at extended intervals, similar to these experiments.

Interval feeding does not work as well for higher feeding rates of low to moderate protein feed (energy feeds). For example, if the producer determines that 7 pounds per day of a 20 percent protein supplement (moderate protein concentration) should be provided, then 49 pounds of feed would need to be delivered each week. With three feedings per week, 16 pounds of feed would have to be delivered at each feeding. With four feedings per week, slightly more than 12 pounds per feeding would need to be provided. These large quantities of feed provided on an interval basis can lead to digestive upset, founder and reduced forage intake and digestibility. Therefore, the maximum recommended amount

Table 4. Performance of beef cows fed supplement three or six times per week.

	Days supplement fed per week	
	3	6
Cow weight in Nov., lb	1,187	1,211
Cow weight loss, Nov. to Apr., lb	242	255
Body condition score, Nov.	5.4	5.4
Body condition score, Apr.	4.4	4.3
Pregnancy rate, %	98	94

Source: Wettemann and Lusby.

to provide during any one feeding event is one percent of body weight (11 pounds for an 1,100-pound cow). Using this rule of thumb and the 1,100-pound cow example, the maximum feed a producer should provide on a daily equivalent would be 4.7 pounds using three feedings per week (11 x 3 = 33 pounds per week; 33 ÷ 7 days per week = 4.7 pounds per day equivalent). The maximum daily equivalent provided using a four-days-per-week schedule would be 6.3 lbs (11 x 4 = 44 pounds per week; 44/7 days per week = 6.3 pounds per day equivalent).

Supplemental Programs for Common Situations in Oklahoma

When hay or pasture nutrient concentration can actually be measured (samples collected and analyzed) and monitored, the methodical approach presented previously will be the most cost effective way to determine the type and amount of supplement to feed. However, many low-cost producers do not feed hay and prefer to use their cows to harvest standing forage. If forage type and conditions are relatively constant from year to year, producers can develop a consistent supplementation program and fine-tune it when necessary. For example, when cattle graze native tallgrass prairie pastures, forage quality consistently declines through the summer, fall and winter months. Protein supplementation needs are quite predictable and may vary more from changing genetics or time of calving than forage conditions. Table 5 shows supplementation schedules for this type of forage with different calving seasons and winter weather conditions. Notice the feeding rate of the high-protein supplement gradually increases to offset the declining forage protein.

More energy is necessary when wet, cold weather conditions persist for long periods of time. Therefore, feeding higher daily amounts of a moderate-protein supplement is advised when these conditions exist or anytime cows are observed to be losing weight and condition too rapidly.

The goal for a spring calving herd is to strive for a BCS 5 in mature cows by the time they calve in order to achieve optimum rebreeding during the spring and early summer months. Fall calving cows usually calve in very good body condition (BCS of 6 to BCS 8) and the producer can allow

these cows to gradually lose some condition through the winter. The main objective for a fall-calving cow is to prevent her from losing too much condition before the end of the breeding season. Once she is pregnant, additional weight and condition loss, and lower rates of supplementation, will not hinder the established pregnancy.

When gestating cows consume hay or pasture that remains above 8 percent protein, low to moderate protein (energy) supplements, such as corn grain, soybean hulls, wheat middlings, or milo can be used at about the same feeding rates as shown in Table 5. However, after calving, a moderate protein supplement may be necessary to offset the protein requirement for lactation. The amount or concentration of protein in the supplement will depend on the protein concentration in the forage base.

Using High Quality Pastures to Supplement Low Quality Forage

In many parts of Oklahoma, small grains pastures can be used to supplement cow herds in winter. Because these are high quality forages, full-time grazing by beef cows results in considerable waste of valuable nutrients. A dry cow grazing continuously on small grain pasture consumes up to 10 times her requirement in protein. A more efficient use of these forages is accomplished by limit-grazing, restricting access to green pasture to a few days or hours each week, and providing low-quality harvested or standing forage during the remaining time.

Small grain forages such as wheat pasture are high in protein, containing 15 percent to 30 percent digestible protein on a dry matter basis. Recent work at the Noble Foundation indicated that mature steers consumed an average of 2.7 pounds of wheat forage dry matter in a 45-minute period. Since the wheat forage contained 30 percent crude protein, the steers consumed 0.8 pound of crude protein during this short period of time. This would be approximately equivalent to 4 pounds of a 20 percent protein supplement. Other research suggested that beef cows consume between 0.5 percent to 1 percent of their body weight in rye forage dry matter during one fill-up grazing bout (Table 6). The fill-up period was approximately four hours in this study. Data suggest small

grains forage dry matter intake is at the lower end of this range during the first few days of limit-grazing. Eventually, small grains forage intake increases substantially during the fill-up grazing bout after the cows have adjusted to the limit-grazing program. After about three weeks, these cows were consuming enough forage to supply about 3 pounds of crude protein; the equivalent of 7.5 pounds of a 40 percent protein supplement or 15 pounds of a 20 percent protein supplement.

Labor availability, location of the small grains pasture, the low-quality forage resource and weather conditions frequently limit the use of limit-grazing systems. For these reasons, producers frequently use an interval limit-grazing approach. Rather than giving cows access to small grains pasture for a few hours each day, cows are provided access to small grains pasture for one fill-up grazing bout (three to five hours) for every two days to six days grazing the low-quality forage or consuming the low-quality harvested forage. A three- to five-hour grazing bout limits the loss of valuable forage due to trampling, bedding down and manure deposits.

The limit-grazing schedule shown in Table 7 is provided as a guideline for limit-grazing intervals necessary to provide adequate supplemental protein and energy to beef cows at different stages of production. For example, in January spring calving, cows would graze native range or consume hay with low protein content for three days, followed by one day (three to five hours) grazing small grains pasture before being returned to the low quality forage source.

Replacement heifers will require approximately one day shorter intervals between small grains grazing bouts to continue growing, maintain or improve body condition and have a reasonable chance of rebreeding for their second calf. Remember that the appropriate time spent grazing the small grains pasture is likely to vary considerably depending on the situation. Factors such as low -quality forage protein and content and digestibility (energy content), small grains forage standing crop, cow size, stage of production, genetic potential for milk production, body condition score and age will have a substantial impact on this decision.

With average weather conditions in central and western Oklahoma, enough small grains forage should be accumulated by early December to supply the protein needs of about 1 cow to 1.5 cows per acre through the middle of February, assuming that a limit-grazing program is used. After the small grains forage begins to grow rapidly during late February or early March, protein needs can be met for 1.5 cows to 3 cows per acre, again, assuming that a limit-grazing program is used.

While not as abundant in protein as small grain forage, tall fescue in winter will meet the protein needs of a dry cow with less than full-time grazing. An efficient system for wintering cows on fescue is to accumulate fall growth in the pasture for grazing after December 1. When pastures are adequately fertilized with nitrogen, the accumulated forage contains from 9 percent to 14 percent protein. Similarly, fertilized, stockpiled Bermudagrass pasture can contain 9 percent to 14 percent protein through the month of December.

Limiting Feed Intake with Salt

Occasionally, it is desirable to self-feed supplements to cows in winter. For example, rough and inaccessible pastures limit a producer's ability to deliver supplements on a timely basis. In these situations, salt can be used to control intake of

Table 6. Beef cow rye forage intake during one fill-up period (approximately 4 hours).

Days relative to initiation of limit-grazing	Forage dry matter intake, lbs	Crudeprotein intake, lbs
Day 1	5.0	1.25
Day 2	7.2	1.80
Day 23	11.9	2.98

Source: Altom and Schmedt.

Table 7. Approximate interval between small grains grazing bouts necessary to meet supplemental protein and energy needs of beef cows^{a,b}.

Month	Number of days consuming low quality forage per fill-up grazing bout	
	Spring calving cows	Fall calving cows ^c
December	4	2
January	3	2
February	3	3
March	2	3
April	2	3
Total days grazing small grains pasture (12/1 – 4/15)	38	42

^a These suggested intervals assume that abundant low quality forage is provided at all times when the cows are not grazing small grains forage.

^b Reduce the suggested interval by one day for first-calf heifers.

^c Calves should be provided free-choice access to the small grains forage using creep gates.

the supplement. The ratio of salt to supplement can be varied to achieve any desired supplement intake.

Self-feeding of supplements tend to allow timid, slow eating cows to get their share. Vitamin A, minerals and other feed additives can be provided through the supplements.

There are disadvantages to feeding salt-concentrate mixes. Salt is not a precise regulator of intake since certain individuals will tolerate more salt than others. Additionally, salt is destructive to metal storage bins, feeders and farm vehicles.

Daily salt requirement for mature cattle is less than 1 ounce per head per day; however, voluntary intake often exceeds minimum needs. Maximum daily voluntary intake of salt is approximately 0.1 pound salt per 100 pounds body weight for most classes of cattle.

Effects of High Salt Intake

Salt toxicity is seldom seen in cattle because of their high tolerance for salt. The one-time lethal dose for mature cattle is 4 to 5 pounds of salt. Salt is rapidly absorbed from the intestinal tract into the bloodstream. It is then excreted

Table 5. Common supplementation strategies for cows grazing native warm-season forage^a during winter.

Month	Spring Calving Cows		Fall Calving Cows	
	Good cow condition and/or moderate weather	Marginal cow condition and/or severe weather	Good to moderate cow condition and/or moderate weather	Thin cow condition and/or severe weather
October	None	None	1 lb HP	1 lb HP
November	1 lb HP ^b	1 lb HP	2 lb HP	2 lb HP
December	2 lb HP	2 lb HP	3 lb HP	3 lb HP
January	3 lb HP	3 lb HP	3 lb HP	6 lb MP
February	3 lb HP	5 lb MP ^c	3 lb HP	7 lb MP
March	3 lb HP	6 lb MP	3 lb HP	7 lb MP
April	2 lb HP	5 lb MP	2 lb HP	6 lb MP

^a Forage protein declines to a low of around 3% to 4% during mid winter.

^b HP = high protein supplement, such as 38% protein range cubes or cotton seed meal.

^c MP = moderate protein supplement, such as 20% protein range cubes or corn gluten feed.