

Lusby, K. S. and R. P. Wettemann. 1980. Effects of early weaning calves from first calf heifers on calf and heifer performance. Okla. Agri. Exp. Stat. MP-107:55.

Wyatt, R. D., R. P. Wettemann, M. B. Gould, Leon Knori, and Robert Totusek. 1976. Effect of single vs twin rearing on cow and calf performance. Okla. Agri. Exp. Sta. MP-96:43.



The Use of Early Weaning in Practical Cattle Management

David Lalman
Extension Beef Cattle Specialist

Introduction

On a short-term basis, many cattlemen find themselves with cows or heifers that are too thin at breeding time to project inadequate rebreeding rates. Droughts, range fires or other natural disasters often leave cattlemen with emergency shortages of forage. Management solutions, even drastic ones that can salvage high pregnancy rates (and therefore, next year's income) are important.

On a longer-term basis, the cattle industry faces severe price competition from poultry and pork. It is unlikely beef cattle will ever be able to compete on a price per pound of product basis with either, especially poultry. Of all the costs of producing beef, the largest single cost is that of investment cost in land for the cow herd, followed by costs for purchased feed and harvested forage. Therefore, management practices reducing nutrient requirements of the cow herd and offering the potential for more efficient utilization of pasture resources need to be evaluated by cattle producers.

Early weaning has been shown to be a possible solution for the short-term forage and reproduction during emergency situations. This practice may also offer some help in the longer-term economic problem if management of the early-weaned calf can be handled.

Early Weaning and Reproduction

The greatest constraints for limiting nutrient intake of beef cows are the necessity of maintaining good body condition for reproduction and increased requirements for milk production. Milk production effects requirements two ways: directly for production of milk, and indirectly in terms of increased maintenance requirements during lactation.

The adverse effects of suckling on reproduction of beef cows has been well researched. Work in the 1930s showed that Holsteins milked six times each day rebred more slowly than cows milked two times or three times (see Table 1). Simulated twinning was produced by grafting an extra calf at birth on Hereford x Holstein cow at the Fort Reno Station (Wyatt et al., 1976). Cows were fed so weight and condition changes were the same for cows with twins and singles. Cows rearing twins produced 39 percent more milk, weaned 60 percent more calf, but were slower to start cycling after calving. At 80 days after calving, only 43 percent of cows with twins had cycled, compared with 71 percent for cows with singles. Cows rearing twin calves were nursed more frequently (4.7 vs 3.4 times daily) and 25 percent longer each day than cows with single calves.

Oklahoma Cooperative Extension Fact Sheets are also available on our website at: <http://osufacts.okstate.edu>

Table 1. Reproduction, performance and nursing behavior of cows rearing single or simulated twin calves.

	<i>Single calves</i>	<i>Twin calves</i>
Daily winter supplement ¹ post-calving, lb	3.9	6.8
Fall weight, lb	1092	1119
Winter weight loss, %	19.0	18.6
Daily milk yield, lb	14.5	20.2
Calving date	12-29-74	12-27-74
Cows cycling by 60 days post-partum, %	36	14
Cows cycling by 80 days post-partum, %	71	43

¹30% all-natural crude protein supplement.

The potential for using early weaning to salvage reproduction is illustrated by an OSU study (Table 2, Lusby and Wettemann, 1980). Calves from 31 spring-calving, 2-year-old first-calf heifers in very thin body condition (condition scores of 3 and 4) were weaned at six to eight weeks of age. Thirty additional heifers were assigned to raise their calves to the normal weaning age of seven months. Early weaned calves were moved to drylot and fed a complete ration made of corn, cottonseed hulls, and protein. At 4 1/2 months of age, half of the early-weaned calves were moved to native pasture with the drylot ration available in a creep feeder. Early weaning improved reproduction rates from 59 percent in heifers that raised their calves to normal weaning up to 97 percent for very thin heifers with early-weaned calves. Weaning weights were similar for drylot and normally weaned heifers (Table 3). However, calves moved to pasture with the drylot ration as a creep feed weighed less.

A second trial looked at the effects of early weaning on reproduction of mature cows in moderate body condition (condition score = 5; Lusby and Parra, 1981). Half the calves from 48 Hereford cows were early weaned at six weeks to eight weeks of age, and the other half remained with their dams until normal weaning age. Early-weaned calves were raised to normal weaning age on the same ration used in the previous trial. Cows with early-weaned calves rebred faster after

Oklahoma State University, in compliance with Title VI and VII of the Civil Rights Act of 1964, Executive Order 11246 as amended, and Title IX of the Education Amendments of 1972 (Higher Education Act), the Americans with Disabilities Act of 1990, and other federal and state laws and regulations, does not discriminate on the basis of race, color, national origin, genetic information, sex, age, sexual orientation, gender identity, religion, disability, or status as a veteran, in any of its policies, practices or procedures. This provision includes, but is not limited to admissions, employment, financial aid, and educational services. The Director of Equal Opportunity, 408 Whitehurst, OSU, Stillwater, OK 74078-1035; Phone 405-744-5371; email: eeo@okstate.edu has been designated to handle inquiries regarding non-discrimination policies: Director of Equal Opportunity. Any person (student, faculty, or staff) who believes that discriminatory practices have been engaged in based on gender may discuss his or her concerns and file informal or formal complaints of possible violations of Title IX with OSU's Title IX Coordinator 405-744-9154.

Issued in furtherance of Cooperative Extension work, acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture, Director of Oklahoma Cooperative Extension Service, Oklahoma State University, Stillwater, Oklahoma. This publication is printed and issued by Oklahoma State University as authorized by the Vice President, Dean, and Director of the Division of Agricultural Sciences and Natural Resources and has been prepared and distributed at a cost of 20 cents per copy. Revised 0715 GH.

Table 2. Weight changes and conception rates of heifers with normal or early weaned calves.

	<i>Suckled</i>	<i>Early Weaned</i>
Heifer weights 11/15/78	738	726
After calving, Feb, Mar, Apr, 1979	698	680
Weight gains changes		
calving to breeding	-16	34
during breeding period (5/29/79-7/31/79)	48	73
calving to weaning (10/11/79)	90	195
Weight at weaning	788	875
Conception rates		
conceived/exposed	19/32	30/31
% pregnant	59.4	96.8

Table 3. Weight gains, weaning weights and feed efficiencies of normal and early weaned calves.

	<i>Suckled</i>	<i>Drylot</i>	<i>Drylot-Pasture</i>
Number of calves	30	16	13
Weight at early weaning, lb		124	126
Gain from 7/31/79-10/11/79		106	62
Weight at normal weaning age	373	374	330
Feed conversion (as fed basis)		4.67	4.78
Feed consumption (lb/hd/day)			
ration 1		2.5	—
ration 2		5.9	4.3
ration 3		11.5	6.5

calving (46 days vs 81 days) and had higher conception rates (100 percent vs. 83 percent; Table 4). Early-weaned calves weighed more at normal weaning age than calves raised by their dams, and gains of early weaned calves were efficient.

Other studies have shown the response to early weaning may depend on body condition score of the cow. Bishop and Wettemann (1990) reported 100 percent of cows with condition scores greater than 5 had begun cycling within 25 days after their calves were early weaned at 45 days of age compared to only 43 percent of cows with condition scores of less than 5. Research has shown that body condition of the cow does influence the time between early weaning and the first postpartum estrous cycle. Cows in moderate to good condition generally cycle within 14 days after calves are weaned. Thinner cows will require more time.

Early Weaning and Forage Intake

A major expense in the production of weaned calves is the cost of forage and the land on which it is grown. Cost of forage can be even more critical during periods of drought when all feed for the cow herd must be purchased. Early weaning can be a good emergency tool during these times. An Illinois study showed that early weaned (110 days of age) fall-calving beef cow-calf pairs consumed 20 percent less total digestible nutrients, fed as hay, than normal weaned pairs. Additionally, the early weaned pairs were 43 percent more efficient in converting TDN to weight gain.

Table 4. Effects of early weaning on cow performance and performance of normal reared and early weaned calves.

	<i>Normal weaned</i>	<i>Early weaned</i>
Cow weight, lb		
Time of early weaning (Avg. date, May 19)	816	832
End of breeding, July 7	922	968
At weaning	920	1040
Condition scores		
Time of early weaning	5.04	5.07
End of breeding	5.69	6.29
At weaning	5.99	6.82
Interval from calving to first observed estrus, days	81	46
No. cows pregnant/exposed	19/23 (83%)	23/23(100%)
Calf weights, lb		
At birth	71	68
At early weaning	145	155
At weaning, (205 day, steer equiv.)	347	435
Avg daily gain, early weaning to 205 days, (158 days)	1.27	1.77
Avg daily feed	—	8.84
Lb feed/lb gain, Dry matter basis	—	4.50

The following information is also discussed in ANSI-3031, Nutrition and Management Considerations for Preconditioning Home Raised Beef Cattle.

What to Do With the Calves

Although the benefits of early weaning on improving reproduction and reducing feed inputs to the cow have been recognized for many years, the factor limiting practical application of early weaning has been management of the early weaned calf. To date, the only programs for spring-born calves that permit "normal" rates of gain to typical normal weaning ages have been drylot programs. Young calves can be very efficient on high-concentrate rations and dry matter conversions of 4:1 are possible up to weights of about 500 lbs. However, feeding programs for young (two- to seven-month-olds) calves need to be "growing programs" that hold daily gains to levels similar to those achieved on the cow. These rates of gain will generally range in the 2.0 lb/day to 2.5 lb/day range, depending on frame size and growth potential of the calves. Otherwise, full-fed baby calves will get fat too early and will not finish at acceptable slaughter weights. Specialty programs, such as limit-feeding of high concentrate rations are attractive to facilities than can manage the programs because of the efficiency and low cost that can be attained.

An "early weaning ration" developed during the OSU studies is shown in Table 5. This is a very palatable ration that is readily consumed by young calves. Intake should be about 3 percent of body weight within seven days. Gains to normal

Table 5. Ration for receiving very light calves or early weaning.

<i>Ingredient</i>	<i>% in ration, as fed</i>
Cottonseed hulls	14.7
Alfalfa pellets	15.1
Rolled corn	46.0
Cane molasses	4.7
Soybean meal	17.7
Calcium carbonate	1.0
Dicalcium phosphate	.5
Salt .3	
Vitamin A	2000 IU/lb
Vitamin E	20 IU/lb
Trace mineral	Cu, Zn, Se as needed to meet NRC req.
Coccidiostat	As per veterinarian's preference
<i>Ration Specs (DM basis)</i>	
Dry matter, %	88.2
NE _m , Mcal/100 lb	82
NE _g , Mcal/100 lb	50
Crude Protein, %	16.7
Potassium, %	1.2
Calcium, %	.95
Phosphorus, %	.45

weaning age will be in the 2.0 lb/day to 2.5 lb/day range. Of course, endless possibilities exist for such rations but this one has been very successful for early weaned calves and stressed light-weight stockers.

As a rule, cattle need to be moved to progressively increasing levels of energy as they move toward slaughter. Moving calves from a high rate of gain in drylot to native range, for example, will result in poor gains. Therefore, calves raised in drylot to normal weaning age on mixed rations need to be moved to nutritional programs with good gain potential after finishing the drylot growing phase. Calves raised to seven months of age in drylot probably need to be moved to a feedlot or a forage with the gain potential of wheat pasture.

Oklahoma has one forage system that may provide an answer for fall-calving programs needing to early wean, namely wheat pasture. In a preliminary trial at OSU, 55 calves born in September and October 1993, were weaned on December 14, held in drylot on prairie hay and 2 lbs of protein pellets, and moved to wheat pasture on December 29. Calves gained about 2 lbs/day on wheat through grazeout with no supplemental feed. Virtually all cows rebred with minimal supplement on native range. It offers the potential for reducing feed costs, increasing stocking rates on the native pasture, and doing this with minimal labor.

Managing Health of Early Weaned Calves

Spring-calving herds present a couple of problems that should not be serious concerns for fall-calving herds. First, there presently is no summer equivalent to wheat pasture for grazing of early weaned calves. Therefore, any early weaning

program for spring-born calves will need to utilize some type of drylot feeding program. Growing programs for early weaned calves studied at OSU have involved full-fed rations (Lusby, et al, 1981; Gill et al., unpublished) and limited program-fed rations (Lusby, et al., 1991).

Minimal health problems have been observed with university and producer cattle demonstrations, when the calves were maintained on the home ranch. However, serious problems have been encountered when early weaned calves were moved directly from the cow to commercial feeding or growing facilities. There must be some kind of prior preparation phase before exposing young, early weaned calves to the stresses of shipment, weaning and the numerous pathogens circulating around a feedlot. The preparation phase will probably involve some concentrate feeding at the ranch of origin and a well-designed immunization program. More studies are needed to involve large numbers of calves from commercial ranches rather than small numbers of Experiment Station calves.

Vaccination programs for 2- to 4-month-old calves are obviously tricky and are left to the discretion of the attending veterinarian. Programs used to date have ranged from only clostridial vaccines at early weaning followed by virus and other vaccines at older ages to aggressive programs with the full gamut of vaccines at early weaning and repeated revaccinations.

Observations and Conclusions

1. Early weaning is a fairly predictable method of salvaging high reproduction rates during droughts, management mistakes and other emergencies.
2. The early weaned calves are pretty tough and learn to eat palatable rations quickly. Health problems when calves are weaned and kept on the ranch have been minimal. Stressful procedures like castration and dehorning should be performed well ahead of early weaning.
3. If at all possible, early weaned calves should be managed on the ranch or managed with only a short haul for some period after weaning.
4. Management of early weaned calves in commercial feedlots is risky and requires close coordination of the rancher, feedlot manager, nutritionist and veterinarian. Each must understand the program and all must cooperate. Mistakes have long-term and often fatal consequences.
5. Wheat pasture offers the potential of high rates of gain for early weaned fall-born calves.
6. There are not presently any forage-based programs for early weaned spring-born calves during the summer that permit gains equal to that on the cow. Crabgrass may offer promise but has not been researched. Research at OSU suggests that gains of about 1.5 lbs/day to 1.75 lbs/day can be achieved with 3 lbs/day or 4 lbs/day of supplemental feed and summer grass. While not a great rate of gain, it is economical and the calves probably would do very well on wheat in the fall.

Literature Cited

- Lusby, K. S. and Donald R. Gill. 1994. Receiving and growing rations. Okla. Coop. Ext. Serv., E-900:7.
- Lusby, K. S. and Angel A. Parra. 1981. Effects of early weaning on calf performance and on reproduction in mature cows. Okla. Agri. Exp. Stat. MP-108:64.