



The Ogallala Aquifer

The Oklahoma Cooperative Extension Service Bringing the University to You!

The Cooperative Extension Service is the largest, most successful informal educational organization in the world. It is a nationwide system funded and guided by a partnership of federal, state, and local governments that delivers information to help people help themselves through the land-grant university system.

Extension carries out programs in the broad categories of agriculture, natural resources and environment; family and consumer sciences; 4-H and other youth; and community resource development. Extension staff members live and work among the people they serve to help stimulate and educate Americans to plan ahead and cope with their problems.

Some characteristics of the Cooperative Extension system are:

- The federal, state, and local governments cooperatively share in its financial support and program direction.
- It is administered by the land-grant university as designated by the state legislature through an Extension director.
- Extension programs are nonpolitical, objective, and research-based information.
- It provides practical, problem-oriented education

for people of all ages. It is designated to take the knowledge of the university to those persons who do not or cannot participate in the formal classroom instruction of the university.

- It utilizes research from university, government, and other sources to help people make their own decisions.
- More than a million volunteers help multiply the impact of the Extension professional staff.
- It dispenses no funds to the public.
- It is not a regulatory agency, but it does inform people of regulations and of their options in meeting them.
- Local programs are developed and carried out in full recognition of national problems and goals.
- The Extension staff educates people through personal contacts, meetings, demonstrations, and the mass media.
- Extension has the built-in flexibility to adjust its programs and subject matter to meet new needs. Activities shift from year to year as citizen groups and Extension workers close to the problems advise changes.

Saleh Taghvaeian

Assistant Professor and Extension Specialist, Water Resources

R. Scott Frazier

Associate Professor and Extension Specialist, Energy Management

Dustin Livingston

Research Assistant

Garey Fox

Professor and Buchanan Endowed Chair, Hydrology and Director, Oklahoma Water Resources Center

The Ogallala aquifer, also known as the High Plains aquifer, is the primary source of water for many communities throughout the High Plains region. Stretching from South Dakota to Texas, the Ogallala aquifer is one of the largest aquifer systems in the U.S. It underlies nearly 122 million acres of land, used primarily for agriculture, producing nearly one-fifth of the nation's wheat, corn and cotton [3, 11]. Approximately 14 percent of the total aquifer area consists of irrigated acres capable of producing \$7 billion in crop sales [1]. The Ogallala aquifer provides one-fourth of the total water supply used for agricultural production across the U.S. [3]

During the 1950s and 1960s, agriculture in the High Plains began to expand both in irrigation systems and irrigated acres [11]. The new irrigation systems provided a way to grow crops even during drought; however, larger and deeper wells were required to meet demands. The irrigation expansion and increased pumping resulted in considerable losses of water storage in parts of the Ogallala aquifer. As a result, Congress directed the U.S. Geological Survey to start monitoring water level changes in this critical groundwater resource, a mission that has continued. The data collected by USGS have yielded critical water-level information, specifically characterizing declines.

The most recent USGS publication estimates a drop in water level of about 15 feet, when averaged over the entire Ogallala area [6]. However, the change in water level is highly variable in different areas of the aquifer; no change or even water level rise was observed mainly in the northern area and significant declines measured in the southern High Plains (Figure 1). When comparing 2013 water levels to those measured in the 1950s, the maximum water rise was 85 feet at an observation well in Nebraska and the maximum water decline was 256 feet at a well in Texas.

The southern High Plains has an arid/semi-arid climate with rainfall amounts less than evaporation and far less than the water requirement of many agricultural crops. The Ogallala

Oklahoma Cooperative Extension Fact Sheets are also available on our website at: <http://osufacts.okstate.edu>




Figure 1. Change in the Ogallala aquifer water level from predevelopment (1950s) to 2013. Data obtained from [6].

aquifer in this region has a very low recharge rate; resulting in very little precipitation reaching the groundwater. Water users in the southern High Plains have experienced significant declines in water levels, and consequently, increases in pumping costs. Areas like the Texas Panhandle and southwestern Kansas have created local districts to monitor water levels and help stakeholders sustain the groundwater supply. In Texas, Groundwater Conservation Districts are formed by people living in geographic areas where groundwater has declined. These GCDs regulate well spacing and pumping rates within their boundaries to ensure long-term groundwater protection and conservation [10].

Oklahoma State University, in compliance with Title VI and VII of the Civil Rights Act of 1964, Executive Order 11246 as amended, and Title IX of the Education Amendments of 1972 (Higher Education Act), the Americans with Disabilities Act of 1990, and other federal and state laws and regulations, does not discriminate on the basis of race, color, national origin, genetic information, sex, age, sexual orientation, gender identity, religion, disability, or status as a veteran, in any of its policies, practices or procedures. This provision includes, but is not limited to admissions, employment, financial aid, and educational services. The Director of Equal Opportunity, 408 Whitehurst, OSU, Stillwater, OK 74078-1035; Phone 405-744-5371; email: eeo@okstate.edu has been designated to handle inquiries regarding non-discrimination policies: Director of Equal Opportunity. Any person (student, faculty, or staff) who believes that discriminatory practices have been engaged in based on gender may discuss his or her concerns and file informal or formal complaints of possible violations of Title IX with OSU's Title IX Coordinator 405-744-9154.

Issued in furtherance of Cooperative Extension work, acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture, Director of Oklahoma Cooperative Extension Service, Oklahoma State University, Stillwater, Oklahoma. This publication is printed and issued by Oklahoma State University as authorized by the Vice President for Agricultural Programs and has been prepared and distributed at a cost of 42 cents per copy. 0616 GH.

Oklahoma

The Ogallala aquifer is one of the major aquifers in Oklahoma, underlying the Panhandle and parts of the northwest regions. In the eastern part of the Oklahoma Panhandle, the Ogallala formation often sits directly on top of consolidated sediments, which are more than 250 million years old. In Cimarron and western Texas counties, a younger formation of shale and sandstone called the Dockum group overlies the redbed. Other formations, including the Dakota Sandstone and Morrison Formation, also are found in the Oklahoma Panhandle [4].

The Panhandle region has a semi-arid climate with the average annual rainfall of about 20 inches during the past 120 years (1895-2015). Drought is common in the area, with one out of every six years having received less than 16 inches of rainfall during the same 120-year period. The Ogallala aquifer supplies more than 98 percent of total water demand in the Oklahoma Panhandle and other sources (e.g. alluvial aquifers and streams) contribute less than 2 percent [9]. Alluvial aquifers, which are shallower and generally connected to surface water resources, are limited. Local rivers and creeks (mainly the Beaver River and the upper Cimarron River) can have periods of low to no flow and are unreliable as a major water supply. Optima Reservoir, the only major Panhandle reservoir, does not sustain water.

Agricultural Water Use

Irrigation is the largest use of water in the Panhandle. Based on 2007 crop mix data, there are approximately 230,000 acres of irrigated land in Cimarron, Texas and Beaver counties, requiring more than 290,000 acre-feet of water per year [9]. This is about 85 percent of the total water demand in the region and is expected to increase to about 306,000 acre-feet by 2060. The major irrigated crops are grain corn and wheat, accounting for about three-fourths of the total irrigated area. According to the USDA National Agricultural Statistics Service, the total harvested area of irrigated corn and wheat in 2012 was more than 106,000 and 65,000 acres, respectively. Grain sorghum ranked lower, with a total irrigated area of about 20,000 acres.

The expansion of irrigated agriculture in Oklahoma Panhandle has been a major driving force for economic development and prosperity of this region. However, this growth has come at the cost of declining non-renewable water resources. Figure 2 demonstrates the cumulative number of irrigation wells drilled in each of the four major Oklahoma counties overlying the Ogallala aquifer.

Since the predevelopment period (prior to 1950), about 3,000 irrigation wells have been drilled into the Ogallala aquifer. The largest number of drilled wells (more than half) were in Texas County, followed by Cimarron, Beaver and Ellis counties. Hence, it is not surprising that the largest decline in groundwater has been experienced in Texas County. According to the Oklahoma Water Resources Board, water levels in the Ogallala aquifer have declined more than 70 feet in Texas County and more than 50 feet in Cimarron County since predevelopment. These large declines can be attributed to the high density of irrigation wells.

The highest rate of adding new wells occurred during the 1960s, with about 107 new wells being drilled every year in three counties of the Panhandle area. The period from 2010




Figure 2. The cumulative number of irrigation wells drilled in each of the four major Oklahoma counties overlying the Ogallala aquifer.

to 2015 was characterized by a major drought in the region. During this five-year period, 226, 95 and 31 new irrigation wells were drilled in Texas, Cimarron and Beaver counties, respectively. The water level declines were more than 13 feet in Texas County, 9 feet in Cimarron County and 4 feet in Beaver County over the same period [8].

Figure 3 provides a closer look at the changes in Ogallala aquifer water levels for Oklahoma. As mentioned before, Texas County shows largest declines in water levels, while parts of the Ellis County experienced a rise in water level. Some of the largest declines in water levels across the entire aquifer have happened in Texas Panhandle just south of the Oklahoma border.

Irrigation Systems

Irrigated agriculture in the Oklahoma Panhandle has changed significantly during the past several decades. Figure 4 depicts two satellite images taken in July of 1976 and July




Figure 3. A closer look at the change in the Ogallala aquifer water level from predevelopment (1950s) to 2013. Data obtained from [6].





Figure 4. Satellite images taken in July 1976 (top) and July 2009 (bottom) show the changes in irrigated areas around Guymon, OK.

of 2009 over the city of Guymon and its surrounding area. Green squares are typically flood (gravity) irrigated fields, while green circles represent center pivot (sprinkler) irrigation. As it can be observed, the majority of irrigated fields were under flood irrigation in 1976, with only a few center-pivot systems present in the satellite image. During the next 33 years, most of irrigated fields switched to center-pivot systems. Compared to flood irrigation, center-pivot systems can apply water in smaller amounts and more uniform patterns, resulting in reduced deep percolation and improved crop yield. In addition, the new center pivot systems have nozzles at a lower elevation, which helps reduce direct evaporation and wind drift losses.

The decline in the Ogallala water level has created numerous challenges for irrigated agriculture in the Oklahoma Panhandle. A main challenge is the reductions in pumping rates, resulting in higher energy costs and decreased revenue,

as crop yields are decreased due to not meeting the full crop water requirement or planting a smaller area. In addition, as the Ogallala level decline and water is being extracted from deeper parts of the aquifer the quality of water may degrade [2].

Since the Ogallala aquifer has a very low recharge rate [7] and other water resources are so scarce, efforts to increase the life of this valuable resource should focus on managing demand. This requires a close collaboration between stakeholders, state/federal agencies and universities. Local groups of stakeholders, such as the Oklahoma Panhandle Agriculture and Irrigation Association, have been active in this field. An example of their efforts is the development of the Panhandle Regional Water Plan in collaboration with other local groups.

References

- Gollehon, Noel and Winston, Bernadette. 2013. Groundwater Irrigation and Water Withdrawals: The Ogallala Aquifer Initiative. *U.S. Geological Service REAP Report*. 2013.
- Hart, D. L., Hoffman, G. L. and Goemaat, R. L. 1976. Geohydrology of the Oklahoma Panhandle, Beaver, Cimarron, and Texas Counties. *U.S Geological Service Report*. 1976.
- Houston, Natalie A., et al. 2013. Geodatabase Compilation of Hydrogeologic, Remote Sensing, and Water-Budget-Component Data for the High Plains Aquifer, 201. *U.S. Geological Service Report*. 2013.
- Luckey, Richard R. and Becker, Mark F. 1999. Hydrogeology, water use, and simulation of flow in the High Plains aquifer in northwestern Oklahoma, southeastern Colorado, southwestern Kansas, northeastern New Mexico, and northwestern Texas. *U.S. Geological Survey Water-Resources Investigations Report* 99-4104. 1999.
- Luckey, Richard R. and Osborn, Noel I. 2000. Water Flow in the High Plains Aquifer in Northwestern Oklahoma. *U.S Geological Service Fact Sheet* 081-00. 2000.
- McGuire, Virginia L. 2014. Water-Level Changes and Change in Water in Storage in the High Plains Aquifer, Predevelopment to 2013 and 2011-2013. *U.S Geological Service Report*. 2014.
- Ochsner, T, Fiebrich, C and Neel, C. 2014. Estimating Groundwater Recharge Using the Oklahoma Mesonet. Stillwater, OK: Oklahoma Water Resources Center, 2014.
- OWRB. 2014. Oklahoma Groundwater Report: Beneficial Use Monitoring Program. 2014.
- Smith, Duane and Associates. 2012. Panhandle Regional Water Plan. Oklahoma Panhandle Agriculture and Irrigation Panhandle Regional Economic Development Coalition, Inc. 2012.
- M. D. Smolen, Aaron Mittelstet, and Bekki Harjo (2012) Whose Water Is It Anyway? Comparing the Water Rights Frameworks of Arkansas, Oklahoma, Texas, New Mexico, Georgia, Alabama, and Florida. Oklahoma Cooperative Extension Service, Publication E-1030.
- Winter, Mary and Foster, Cliff. 2014. Ogallala Aquifer-Lifblood of the High plains Part 1: Withdrawals Exceed Recharge. 2014.