



The Oklahoma Cooperative Extension Service
Bringing the University to You!

The Cooperative Extension Service is the largest, most successful informal educational organization in the world. It is a nationwide system funded and guided by a partnership of federal, state, and local governments that delivers information to help people help themselves through the land-grant university system.

Extension carries out programs in the broad categories of agriculture, natural resources and environment; family and consumer sciences; 4-H and other youth; and community resource development. Extension staff members live and work among the people they serve to help stimulate and educate Americans to plan ahead and cope with their problems.

Some characteristics of the Cooperative Extension system are:

- The federal, state, and local governments cooperatively share in its financial support and program direction.
- It is administered by the land-grant university as designated by the state legislature through an Extension director.
- Extension programs are nonpolitical, objective, and research-based information.

- It provides practical, problem-oriented education for people of all ages. It is designated to take the knowledge of the university to those persons who do not or cannot participate in the formal classroom instruction of the university.
- It utilizes research from university, government, and other sources to help people make their own decisions.
- More than a million volunteers help multiply the impact of the Extension professional staff.
- It dispenses no funds to the public.
- It is not a regulatory agency, but it does inform people of regulations and of their options in meeting them.
- Local programs are developed and carried out in full recognition of national problems and goals.
- The Extension staff educates people through personal contacts, meetings, demonstrations, and the mass media.
- Extension has the built-in flexibility to adjust its programs and subject matter to meet new needs. Activities shift from year to year as citizen groups and Extension workers close to the problems advise changes.

Oklahoma State University, as an equal opportunity employer, complies with all applicable federal and state laws regarding non-discrimination and affirmative action. Oklahoma State University is committed to a policy of equal opportunity for all individuals and does not discriminate based on race, religion, age, sex, color, national origin, marital status, sexual orientation, gender identity/ expression, disability, or veteran status with regard to employment, educational programs and activities, and/or admissions. For more information, visit <https://eeo.okstate.edu>.

Issued in furtherance of Cooperative Extension work, acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture, Director of Oklahoma Cooperative Extension Service, Oklahoma State University, Stillwater, Oklahoma. This publication is printed and issued by Oklahoma State University as authorized by the Vice President for Agricultural Programs and has been prepared and distributed at a cost of 20 cents per copy. 04/2020 GH.

EXTENSION

Oklahoma Cooperative Extension Fact Sheets are also available on our website at: extension.okstate.edu

Oklahoma Bermudagrass Variety Performance Tests: 2019 Forage Year

Alex Rocateli
 Forage Systems Extension Specialist

Lucas Freires Abreu
 Graduate Research Assistant, Forage Systems

Introduction

Selecting the most appropriate variety is very important for establishing perennial pastures, such as Bermudagrass, because variety selection will dictate the potential forage yield and quality for many years to come. Establishment method (sprigging vs. seeding), winter hardiness, site-specific adaptations, growth morphology (grazing vs. hay types) and forage quality are all traits that must be considered when deciding which Bermudagrass variety to plant. Fact sheet PSS-2600, *Selecting an Appropriate Bermudagrass Variety for Pastures*, available at: extension.okstate.edu, discusses the traits that are important for choosing the most appropriate variety for a specific location.

The purpose of this current report is to highlight differences in forage yield among commercially available sprigged

and seeded Bermudagrass varieties grown in trials in several locations across Oklahoma. This information can give producers an indication of forage potential of the varieties evaluated. However, the forage yields reported here are the results derived from complex interaction among varietal genetic potential, management factors such as planting date, seeding rate, soil fertility and environmental factors, such as temperature, rainfall amount and distribution. These factors vary greatly from location to location and among Oklahoma's geographic regions, so final forage yield and quality observed by producers will be affected by environmental and management. Thirteen commercially available varieties were tested in four different locations across Oklahoma (Figure 1). Fact sheet PSS-2600 contains brief descriptions of the agronomic characteristics of all tested varieties.




Figure 1. Bermudagrass performance test locations.

Table 1. Site Description and Methods.

Locations	Establishment method	Establishment date	Soil texture	pH	N	STP	STK
Perkins	Sprigged	7/15/16	sandy loam	6.3	0	17	110
Chickasha	Sprigged	7/18/16	clay loam	6.6	3	67	510
Chickasha	Seeded	5/6/18	silt loam	6.4	9	76	336
Haskell	Sprigged/seeded	5/18/17	silt loam	7.1	4	42	103
Lane	Sprigged	5/20/17	sandy loam	6.9	5	18	122

STP: soil test P index; STK: soil test K index

Site Description and Methods

Bermudagrass variety performance tests were established at four locations in Oklahoma in 2016 and 2017. Forage yields were harvested in successive years. The experimental design was a randomized complete block with four replications. Plots were established in a conventionally-tilled, firm seedbed. Phosphate (P) and potash (K) were applied according to soil test recommendations (Table 1) and incorporated 3 inches deep during seedbed preparation. Sprigged plots were sprigged at four sprigs per 10 square feet, which is similar to a rate of 40 to 50 bushels of sprigs per acre. Seeded plots were seeded at 15 pounds of pure live seed per acre, which corresponds to the highest recommended seeding rate among participating companies. Thirty pounds of actual nitrogen per acre (65 pounds of urea per acre) was broadcast during establishment. Nitrogen fertilizer was broadcast at a rate of 200 pounds nitrogen per acre (440 pounds of urea per acre) right after Bermudagrass broke dormancy and produced the first green leaves (green-up) every year.

Plots were hand-weeded during the establishment phase to avoid potential herbicide injury. After establishment, pre-emergent (pendimethalin) and post-emergent (glyphosate) herbicides were applied in late winter or early spring (before green-up), respectively. Early-season weeds at green up (early May) were controlled by mowing at 2 to 3 inches in height. In-season weeds were controlled by hand-weeding and/or selective post-emergent herbicides such as 2,4-D and dicamba.

Forage was measured by hand clipping three 1-square-foot quadrats per plot 1 inch above the soil surface. Plots were mowed and cleaned to simulate hay harvest immediately after sample collection. Samples were oven-dried until constant weight indicating that all samples were completely dry (absent of moisture), and total dry forage yield was assessed. Forage sampling (cut) was targeted for every six weeks, which generally resulted in three to four cuttings per year, depending on temperature and rainfall amount and distribution through the season.

Results

Perkins

Forage yields were similar to those reported in 2018; however, yields were less than in 2017. Herbicides failed to control weeds during the early stages. Abnormal large rainfall

totals in March and April delayed pre-emergent application and prohibited post-emergence herbicide application before Bermudagrass green-up. Consequently, weed interference negatively impacted the first cut yields. Inadequate rainfall after first cut consequently limited second cut yields. The third cut was the least productive of the season; however, it was similar to previous years.

Chickasha

Large amounts of rainfall during early spring limited the effect of the pre-emergent herbicide and prohibited the application of a non-selective post-emergent herbicide before green-up. As a result, weed interference was substantial during the early season, which decreased forage yield when compared to the previous year. Inadequate rainfall after the first cut reduced forage yield in the second cut. As expected, the third cut was the least of the season; however, it was similar to the last years.

Among sprigged cultivars, total forage yields ranged from 7.0 to 3.9 tons per acre. The top highest-yielding varieties were Ozark, Tipton44 and Goodwell, respectively. Greenfield was the lowest yielding variety.

Good total forage production was observed among seeded cultivars at their first-year production. All seeded cultivars showed similar total dry forage yields ranging from 5.7 to 5.0 tons per acre.

Haskell

Although amounts of rainfall during early spring were greater than optimal in Haskell, pre- and post-herbicides were applied during early spring; therefore, weed incidence was much less than at other locations. However, a large number of cloudy days during the early season reduced air temperature, total solar radiation, and consequently, yields at first cut when compared to the previous year. Second and third cuts yields were similar to the previous year. No weeds or water stress were observed throughout the entire growing season.

Ozark (6.0 tons per acre) followed by Midland 99 and Goodwell were the top total forage yielding sprigged cultivars, and Greenfield (3.9 tons per acre) was the lowest yielding cultivar. Among seeded cultivars, the top total yielding varieties were Riata (6.0 tons per acre), Stampede Plus and Wrangler. Tierra Verde (4.8) was the lowest yielding.

Lane

Much like Perkins and Chickasha, delayed pre- and post-herbicide application resulted in weed interference during




Figure 2. Average daily temperature (line graph) and rainfall (bar graph) from April 15 to October 15 in each respective location and year. Weather data courtesy of the Oklahoma Mesonet.

Forage Yield Summary - Past Years

Entry	Perkins		Chickasha		Haskell	Lane
	2017	2018	2017	2018	2018	2018
	<i>Tons Dry Matter/Acre</i>					
Sprigged						
Goodwell	7.4	5.0	7.7	8.1	6.2	6.0
Ozark	7.0	5.0	6.1	9.2	7.6	7.4
Midland99	6.6	5.0	6.3	9.8	6.6	7.3
Midland	6.3	4.7	5.5	8.2	6.2	6.6
Tifton44	6.2	4.8	5.0	8.9	6.2	7.1
Greenfield	4.1	4.3	6.7	6.6	5.0	5.2
Mean	6.3	4.8	6.2	8.5	6.3	6.6
5% LSD	1.3	NS	NS	2.7	2.4	1.4
Seeded						
Cheyenne II	-	-	-	-	4.7	-
Mohawk	-	-	-	-	4.7	-
Riata	-	-	-	-	7.0	-
Sahara II	-	-	-	-	4.6	-
Stampede Plus	-	-	-	-	6.7	-
Tierra Verde	-	-	-	-	5.9	-
Wrangler	-	-	-	-	4.4	-
Mean					5.4	
5% LSD					1.5	

Design: Randomized Complete Block
 Plot size: sprigged 17 x 17 ft.
 No. of Reps: 4
 Seeded 8 x 17 ft.
 NS= not significant at p = 0.05

Data provided by the Plant & Soil Sciences Department of the Division of Agricultural Sciences and Natural Resources of Oklahoma State University. For additional information, contact Alex Rocateli at alex.rocateli@okstate.edu
 FOR A DETAILED DESCRIPTION OF PAST YEARS ACCESS CR-2605 at <https://extension.okstate.edu/>

the early season. Furthermore, highfield humid conditions resulting in abundance of free moisture in the plant canopy during May and June resulted in a very high incidence of fungal disease, i.e., Bipolaris Leaf Spot (*Bipolaris* spp.) in all cultivars. This disease persisted after the first cutting, affecting plants regrowth. Consequently, the first and second cuts had reduced forage yields when compared to the previous year's yields. All cultivars provided similar total forage yields ranging from 5.9 to 5.0 tons per acre, except for Greenfield (4.4 tons per acre).

Final Considerations

Differences among varieties across locations were observed; however, it is too early to conclude which varieties are best adapted to each location. Bermudagrass takes three to

Sprigged Bermudagrass Performance test, Perkins, 2019

Cimarron Valley Research Station, Payne County

Entry	06/27	08/13	09/30	Total
	<i>Tons Dry Matter/Acre</i>			
Goodwell	2.7	1.2	1.1	4.9
Ozark	2.8	1.0	1.0	4.8
Midland99	2.0	1.3	1.3	4.6
Midland	2.1	1.0	1.0	4.1
Tifton44	2.8	1.2	1.2	5.2
Greenfield	2.5	1.3	0.9	4.7
Mean	2.5	1.2	1.1	4.7
5% LSD	NS	0.3	0.3	NS

Design: Randomized Complete Block
 Plot size: 17 x 17 ft. planted
 No. of Reps: 4
 NS = not significant at p = 0.05

Data provided by the Plant & Soil Sciences Department of the Division of Agricultural Sciences and Natural Resources of Oklahoma State University. For additional information, contact Alex Rocateli at alex.rocateli@okstate.edu

four years to establish and to provide data that are indicative of final yield potential. These Bermudagrass variety performance tests will be continued to assist producers in making future variety decisions.

Acknowledgments

Thanks to the Oklahoma Agricultural Experiment Station (OAES) superintendents Josh Massey, Michael Pettijohn, Rodney Farris and Jim Vaughan for conducting field management practices such as soil bed preparation, fertilization, weed/pest control and alleys maintenance. This research was developed in collaboration with OAES, Oklahoma Cooperative Extension Service (OCES); and partially funded by Pennington® seeds and Johnston Seeds Co.

Sprigged Bermudagrass Performance test, Chickasha, 2019

South Central Research Station, Grady County

Entry	06/14	08/08	09/23	Total
	<i>Tons Dry Matter/Acre</i>			
Goodwell	4.0	1.8	1.1	6.9
Ozark	4.0	2.2	1.2	7.4
Midland99	4.1	2.3	1.0	7.4
Midland	3.5	2.0	1.1	6.6
Tifton44	3.8	2.0	1.1	6.9
Greenfield	3.9	1.3	0.8	5.9
Mean	3.9	1.9	1.1	6.9
5% LSD	NS	0.4	0.3	0.8

Design: Randomized Complete Block
 Plot size: 17 x 17 ft. planted
 No. of Reps: 4
 NS = not significant at p = 0.05

Data provided by the Plant & Soil Sciences Department of the Division of Agricultural Sciences and Natural Resources of Oklahoma State University. For additional information, contact Alex Rocateli at alex.rocateli@okstate.edu

Seeded Bermudagrass Performance test, Chickasha, 2019
South Central Research Station, Grady County

Entry	Licensee	First year production Tons Dry Matter/Acre			
		06/14	08/08	09/23	Total
Cheyenne II	Pennington ®	2.0	1.8	1.2	5.0
Mohawk	Pennington ®	2.1	1.7	1.2	5.0
Riata	Johnston Seed Co.	2.3	1.6	1.3	5.2
Sahara II	Pennington ®	2.4	2.0	1.3	5.7
Stampede Plus	Johnston Seed Co.	2.5	1.9	1.2	5.6
Tierra Verde	Pennington ®	2.7	1.7	1.2	5.7
Wrangler	Johnston Seed Co.	2.6	1.5	1.4	5.5
Mean		2.3	1.8	1.2	5.3
5% LSD		0.7	NS	0.4	NS

Design: Randomized Complete Block
Plot size: 8 x 17 ft. planted
No. of Reps: 4
NS= not significant at p = 0.05

Data provided by the Plant & Soil Sciences Department of the Division of Agricultural Sciences and Natural Resources of Oklahoma State University. For additional information, contact Alex Rocateli at alex.rocateli@okstate.edu

Seeded Bermudagrass Performance test, Haskell, 2019
Eastern Research Station, Muskogee County

Entry	Licensee	06/27	08/13	09/30	Total
Cheyenne II	Pennington ®	2.4	1.6	1.0	4.9
Mohawk	Pennington ®	2.1	1.2	1.2	4.5
Riata	Johnston Seed Co.	2.6	1.8	1.6	6.0
Sahara II	Pennington ®	2.3	1.5	1.0	4.8
Stampede Plus	Johnston Seed Co.	3.0	1.6	1.3	6.0
Tierra Verde	Pennington ®	2.5	1.4	0.9	4.8
Wrangler	Johnston Seed Co.	3.0	1.3	1.0	5.3
Mean		2.6	1.5	1.4	5.2
5% LSD		0.5	0.5	0.3	0.8

Design: Randomized Complete Block
Plot size: 8 x 17 ft. planted
No. of Reps: 4
NS= not significant at p = 0.05

Data provided by the Plant & Soil Sciences Department of the Division of Agricultural Sciences and Natural Resources of Oklahoma State University. For additional information, contact Alex Rocateli at alex.rocateli@okstate.edu

Sprigged Bermudagrass Performance test, Haskell, 2019
Eastern Research Station, Muskogee County

Entry	06/18	08/01	09/25	Total
Tons Dry Matter/Acre				
Goodwell	3.3	1.7	1.0	6.0
Ozark	3.9	2.0	1.1	7.0
Midland99	3.0	1.8	1.1	6.6
Midland	3.4	2.1	1.2	5.7
Tifton44	2.8	1.3	1.1	5.2
Greenfield	2.2	0.9	0.9	3.9
Mean	3.1	1.6	1.1	5.7
5% LSD	0.8	0.4	0.2	1.2

Design: Randomized Complete Block
Plot size: 8 x 17 ft. planted
No. of Reps: 4
NS= not significant at p = 0.05

Data provided by the Plant & Soil Sciences Department of the Division of Agricultural Sciences and Natural Resources of Oklahoma State University. For additional information, contact Alex Rocateli at alex.rocateli@okstate.edu

Sprigged Bermudagrass Performance test, Lane, 2019
Wes Watkins Research and Extension Center, Atoka County

Entry	06/28	08/14	09/30	Total
Tons Dry Matter/Acre				
Goodwell	2.3	1.4	1.3	5.1
Ozark	2.8	2.0	1.1	5.9
Midland99	2.9	1.4	1.3	5.6
Midland	2.6	1.3	1.4	5.3
Tifton44	2.5	1.4	1.2	5.0
Greenfield	2.6	1.0	0.8	4.4
Mean	2.6	1.4	1.2	5.2
5% LSD	0.4	0.4	0.4	0.9

Design: Randomized Complete Block
Plot size: 8 x 17 ft. planted
No. of Reps: 4
NS= not significant at p = 0.05

Data provided by the Plant & Soil Sciences Department of the Division of Agricultural Sciences and Natural Resources of Oklahoma State University. For additional information, contact Alex Rocateli at alex.rocateli@okstate.edu