



Current Report

EXTENSION

Oklahoma Cooperative Extension Fact Sheets are also available on our website at:
extension.okstate.edu

The Oklahoma Cooperative Extension Service WE ARE OKLAHOMA

The Cooperative Extension Service is the largest, most successful informal educational organization in the world. It is a nationwide system funded and guided by a partnership of federal, state, and local governments that delivers information to help people help themselves through the land-grant university system.

Extension carries out programs in the broad categories of agriculture, natural resources and environment; family and consumer sciences; 4-H and other youth; and community resource development. Extension staff members live and work among the people they serve to help stimulate and educate Americans to plan ahead and cope with their problems.

Some characteristics of the Cooperative Extension system are:

- The federal, state, and local governments cooperatively share in its financial support and program direction.
- It is administered by the land-grant university as designated by the state legislature through an Extension director.
- Extension programs are nonpolitical, objective, and research-based information.
- It provides practical, problem-oriented education for people of all ages. It is designated to take the knowledge of the university to those persons who do not or cannot participate in the formal classroom instruction of the university.
- It utilizes research from university, government, and other sources to help people make their own decisions.
- More than a million volunteers help multiply the impact of the Extension professional staff.
- It dispenses no funds to the public.
- It is not a regulatory agency, but it does inform people of regulations and of their options in meeting them.
- Local programs are developed and carried out in full recognition of national problems and goals.
- The Extension staff educates people through personal contacts, meetings, demonstrations, and the mass media.
- Extension has the built-in flexibility to adjust its programs and subject matter to meet new needs. Activities shift from year to year as citizen groups and Extension workers close to the problems advise changes.

Management of Insect and Mite Pests in Corn

Tom A. Royer
Extension Entomologist

Arthropod pests of corn are varied, and often difficult to manage. Many corn pest problems can be avoided by implementing an Integrated Pest Management (IPM) plan that includes preventive pest management practices, such as selecting varieties adapted to Oklahoma growing conditions, planting at an optimal date, proper fertilization and irrigation, and using crop rotations. The application of insecticides, while sometimes necessary, should not be used as a substitute for good agronomic practices or as "preventative insurance" because it is rarely economically or environmentally justifiable.

The information herein is for educational purposes only. Reference to commercial products or trade names is made with the understanding that no discrimination is intended and no endorsement by the Cooperative Extension Service is implied.

Pesticide recommendations in this publication were correct as of the "Modified Date" but always check the label that came with the purchased insecticide for the most current rates and restrictions.

The first name listed is the trade name of a product registered for use in corn for the listed pest. The name in

(parentheses) listed below the trade name is the name of the active ingredient. The active ingredient name is provided because in many cases, there are other registered products containing the same active ingredient that may cost less, so producers should compare prices.

The number [in brackets] following a product is its Mode of Action number [MOA]. The more frequently insecticides with the same MOA are used, the more likely resistance will occur. This number provides an easy way to select different modes of action to avoid selecting for pests that are resistant to a certain mode of action.

Refer to the following publications for additional information on corn pest management.

- AGEC-203 Estimating Yield and Economic Returns from Replanting Corn
- CR-2105 National Corn Handbook: Aflatoxins and other Mycotoxins
- EPP-7160 Field Key to Larvae in Corn
- EPP-7196 Grasshopper Management in Rangeland, Pastures and Crops

The pesticide information presented in this publication was current with federal and state regulations at the time of printing. The user is responsible for determining that the intended use is consistent with the label of the product being used. Use pesticides safely. Read and follow label directions. The information given herein is for educational purposes only. Reference to commercial products or trade names is made with the understanding that no discrimination is intended and no endorsement by the Cooperative Extension Service is implied.

Oklahoma State University, as an equal opportunity employer, complies with all applicable federal and state laws regarding non-discrimination and affirmative action. Oklahoma State University is committed to a policy of equal opportunity for all individuals and does not discriminate based on race, religion, age, sex, color, national origin, marital status, sexual orientation, gender identity/expression, disability, or veteran status with regard to employment, educational programs and activities, and/or admissions. For more information, visit <https://eeo.okstate.edu>.

Issued in furtherance of Cooperative Extension work, acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture, Director of Oklahoma Cooperative Extension Service, Oklahoma State University, Stillwater, Oklahoma. This publication is printed and issued by Oklahoma State University as authorized by the Vice President for Agricultural Programs and has been prepared and distributed at a cost of 40 cents per copy. Revised 04/2021 GH.

Management of Insect and Mite Pests in Corn

<i>Pest, Damage and Treatment Threshold</i>	<i>Insecticide Formulation and [Group]* and (active ingredient)</i>	<i>Rate of Product per Acre or 1,000 ft-row</i>	<i>Comments</i>
<p>Armyworm</p> <p>1 inch to 1.5 inches. Dark green or brown caterpillar with five stripes along a smooth body. Head with honeycomb-like markings.</p> <p>Damage: Armyworms present throughout growing season, but natural enemies have large impact on them.</p> <p>Threshold: Treat if 30% of plants (seedling to six extended leaves) are infested, or when 75% of plants are infested with one or more larvae on larger plants.</p>	Asana XL [3] (esfenvalerate)	5.8 to 9.6 fl oz (0.03 to 0.05 lb.)	21-day waiting period.
	<i>Bacillus thuringiensis</i> [11] (Biobit, Condor, Dipel, Lepinox, Javelin, Xentari)	See product label for specific rates	Check label for waiting periods.
	Baythroid XL [3] (beta-cyfluthrin)	1.6 to 2.8 fl oz (0.013 to 0.022 lb.)	21-day wait for grain or fodder, 0 days for green forage.
	Besiege[28,3] (chlorantraniliprole + lambda-cyhalothrin)	6.0 to 10.0 fl oz/A	21-day wait for harvest.
	Blackhawk [5] (spinosad)	1.67 to 3.3 fl oz (0.038 to 0.075 lb.)	1-day PHI for grain, 7-day wait for grazing, 28 days for fodder.
	Cobalt [1B, 3] (chlorpyrifos + gamma-cyhalothrin)	13 to 26 fl oz	21-day wait for harvest, 1 day for grazing or silage.
	Coragen [28] (chlorantraniliprole)	3.5 to 7.5 fl oz (0.045 to 0.098 lb.)	14-day waiting period for harvest 1 day for forage, silage, stover.
	Delta Gold [3] (deltamethrin)	1.5 to 1.9 fl oz (0.018 to 0.022 lb.)	21-day wait for harvest, 14 days for grazing or silage.
	Elevest [3A,28] (bifenthrin + chlorantraniliprole)	5.6 to 9.6 fl oz (0.098 to 0.167 lb.)	30-day waiting period.
	Fastac [3] (alpha-cypermethrin)	3.2 to 3.8 fl oz (0.020 to 0.025 lb.)	30-day wait for grain, 60 days for silage.
	Hero [3] (zeta-cypermethrin + bifenthrin)	4.0 to 10.3 fl oz	0-day wait for green forage, 21 days for harvest or fodder.
	Intrepid 2F [18] (methoxyfenozide)	1.28 to 1.92 fl oz (0.02 to 0.03 lb.)	3-day wait for forage, 21 days for harvest or grazing.
	Lannate LV [1A] (methomyl)	0.75 to 1.5 pt (0.225 to 0.45 lb.)	35-day waiting period.
	Lorsban 4E [1B] (chlorpyrifos)	1 to 2 pt (0.5 to 1.0 lb.)	30-day wait for grain, 60 days for silage.
	Match-Up [1b,3] (chlorpyrifos + bifenthrin)	5.5 to 16.4 fl oz	30-day wait for grain or harvest.
Mustang MAXX EC [3] (zeta-cypermethrin)	3.2 to 4.0 fl oz (0.02 to 0.025 lb.)	14-day waiting for harvest.	
Pounce 25WP [3] (permethrin)	6.4 to 9.6 oz (0.1 to 0.2 lb.)	30-day waiting period.	
Proaxis 0.5 SC [3] (gamma-cyhalothrin)	2.56 to 3.84 fl oz (0.01 to 0.015 lb.)	21-day waiting period.	
Radiant SC [5] (spinetoram)	3.0 to 6.0 fl oz (0.023 to 0.047 lb.)	28-day wait for harvest, 3 days for forage or fodder.	
Sevin XLR [1A] (carbaryl)	1 to 2 qt (1.5 to 2 lb.)	48-day waiting period for harvest, 14 days for grazing.	
Stallion [1B,3] (chlorpyrifos + zeta-cypermethrin)	9.25 to 11.75 fl oz	30-day wait for harvest, 60 days for grazing or silage.	
Warrior II with Zeon [3] (lambda-cyhalothrin)	1.28 to 1.92 fl oz (0.02 to 0.03 lb.)	21-day wait for harvest or grazing.	

<i>Pest, Damage and Treatment Threshold</i>	<i>Insecticide Formulation and [Group]* and (active ingredient)</i>	<i>Rate of Product per Acre or 1,000 ft-row</i>	<i>Comments</i>
<p>Chinch bug</p> <p>Nymphs are bright red with white band across back. Adults ½ inch, black with white “hour glass” shape on back.</p> <p>Damage: Adults may fly into field, early (March April) or adults and nymphs move in to corn from maturing wheat fields (April-May). Remove plant juices, cause stunting, wilting and reddening of leaves.</p> <p>Threshold:</p> <p>Plants are less than 6 inches: Two or more chinch bugs on 20% of plants</p> <p>Plants are 6 to 18 inches: 10 or more chinch bugs on 75% of plants.</p>	Seed Treatments		
	Cruiser 5FS [4A] (thiamethoxam)	1.13 to 3.61 fl oz / 80,000 seed	Do not feed treated seed. Generally must order through a seed dealer.
	Gaucho 600 [4A] (imidacloprid)	2.7 to 6.0 fl oz/ 80,000 seed	Do not feed treated seed. See label for mixing and handling instructions. Follow all label restrictions.
	Poncho 600 [4A] (clothianidin)	1.13 to 2.26 fl oz/ 80,000 seed	Do not feed treated seed. See label for mixing and handling instructions. Follow all label restrictions.
	Planting Time Applications		
	Force CS [3] (tefluthrin)	0.46 to 0.57 fl oz/ 1,000 ft row	T-band application. Read label carefully for restrictions.
	Post-emergence Sprays		
	Asana XL [3] (esfenvalerate)	5.8 to 9.6 fl oz (0.03 to 0.05 lb ai/A)	21-day waiting period.
	Baythroid XL [3] (beta-cyfluthrin)	1.6 to 2.8 fl oz (0.013 to 0.022 lb ai/A)	21-day wait for grain or fodder, 0 day for green forage.
	Brigade 2EC [3] (bifenthrin)	2.1 to 6.4 fl oz (0.033 to 0.1 lb ai/A)	30 -day waiting period for grain or grazing.
	Cobalt [1B, 3] (chlorpyrifos + gamma-cyhalothrin)	19 to 38 fl oz	21-day wait for harvest.
	Delta Gold [3] (deltamethrin)	1.5 to 1.9 fl oz (0.018 to 0.022 lb ai/A)	21-day wait for harvest, 12 days for grazing or silage.
	Fastac [3] (alpha-cypermethrin)	3.2 to 3.8 fl oz (0.02 to 0.025 lb ai/A)	30-day wait for harvest, 60 days for grazing.
	Hero [3] (zeta-cypermethrin + bifenthrin)	4.0 to 10.3 fl oz	30-day PHI for grain and stover, 60 days for forage.
	Lorsban 4E [1B] (chlorpyrifos)	1 to 2 pt (0.5 to 1.0 lb ai/A)	21-day PHI for harvest.
Match-Up [1B,3] (chlorpyrifos + bifenthrin)	5.5 to 16.4 fl oz	30-day wait for grain or harvest.	
Mustang MAXX EC [3] (zeta-cypermethrin)	3.2 to 4.0 fl oz (0.02 to 0.025 lb ai/A)	7-day waiting period for harvest.	
Proaxis 0.5 SC [3] (gamma-cyhalothrin)	3.84 fl oz (0.015 lb ai/A)	21-day waiting period.	
Sevin XLR [1A] (carbaryl)	1 to 2 qt (0.5 to 1 lb ai/A)	48-day waiting period for harvest, 14 days for grazing or silage.	
Stallion [1B,3] (chlorpyrifos + zeta-cypermethrin)	9.25 to 11.75 fl oz	30-day wait for harvest, 60 days for grazing or silage.	
Warrior II w Zeon [3] (lambda-cyhalothrin)	1.92 fl oz (0.03 lb ai/A)	21-day wait for harvest. Check label for grazing restrictions.	

<i>Pest, Damage and Treatment Threshold</i>	<i>Insecticide Formulation and [Group]* and (active ingredient)</i>	<i>Rate of Product per Acre or 1,000 ft-row</i>	<i>Comments</i>
Corn Earworm Striped robust caterpillars that range in color from green to pink to brown to black. Damage: Caterpillars injure ear tips, feed in whorls. Feeding damage may increase potential for aflatoxins in grain. Threshold: Not practical to control in field corn	Many Bt corn hybrids offer some suppression of corn earworm, but it is not recommended that corn earworm be controlled with insecticides.	NA	
Corn rootworm (adults) Small beetle, with black stripes, 12 spots or green Damage: Feed on silks. Heavy populations may interfere with pollination. Threshold: Treat if beetles are abundant (more than five per plant and silks are being severely clipped).	Asana XL [3] (esfenvalerate) Baythroid XL [3] (beta-cyfluthrin) Besiege[28,3] (chlorantraniliprole + lambda-cyhalothrin) Brigade 2EC [3] (bifenthrin) Cobalt [1B, 3] (chlorpyrifos + gamma-cyhalothrin) Delta Gold [3] (deltamethrin) Dimethoate 4E [1B] Fastac [3] (alpha-cypermethrin) Hero [3] (zeta-cypermethrin + bifenthrin) Lorsban 4E [1B] (chlorpyrifos) Match-Up [1B,3] (chlorpyrifos + bifenthrin) Mustang MAXX EC [3] (zeta-cypermethrin) Pounce 25WP [3] (permethrin) Proaxis 0.5 SC [3] (gamma-cyhalothrin) Sevin XLR [1A] (carbaryl) Stallion[1B,3] (chlorpyrifos + zeta-cypermethrin) Warrior II w Zeon [3] (lambda-cyhalothrin)	5.8 to 9.6 fl oz (0.03 to 0.05 lb ai/A) 1.6 to 2.8 fl oz (0.013 to 0.022 lb ai/A) 6.0 to 10 fl oz 2.1 to 6.4 fl oz (0.033 to 0.1 lb ai/A) 13 to 26 fl oz 1.5 to 1.9 fl oz (0.018 to 0.022 lb ai/A) 0.66 to 1 pt 2.7 to 3.8 fl oz (0.017 to 0.025 lb ai/A) 4.0 to 10.3 fl oz 1 to 2 pt (0.5 to 1 lb ai/A) 5.5 to 16.4 fl oz 3.2 to 4.0 fl oz (0.02 to 0.025 lb ai/A) 6.4 to 9.6 oz (0.1 to 0.15 lb ai/A) 2.56 to 3.84 fl oz (0.01 to 0.015 lb ai/A) 1 to 2 qt (0.5 to 1 lb ai/A) 9.25 to 11.75 fl oz 1.28 to 1.92 fl oz (0.02 to 0.03 lb ai/A)	21-day waiting period. 21-day wait for grain or fodder, 0 day for green forage. 21-day wait for harvest. 30-day wait for grazing or harvest. 21-day wait for harvest, 14 days for grazing or silage. 21-day wait for harvest, 12 days for grazing or silage. 14-day waiting period. 30-day wait for grain, 60 days for silage. 30-day PHI for grain and stover, 60 days for forage. 21-day waiting period. 30-day wait for grain. 7-day waiting period for harvest. 30-day waiting period. 21-day waiting period for harvest or grazing. 48-day waiting period for harvest, 14 days for grazing. 30-day waiting period for grain, 60 days for forage. 21-days for harvest. Check label for grazing restrictions.

Pre-harvest Intervals and grazing restrictions		
Asana XL Aztec 2.1G Baythroid XL Besiege Blackhawk Brigade 2EC Cobalt Comite II Counter 15G	21-day PHI for harvest or grazing Do not exceed 7.3 lb. per acre per crop season 21-day waiting period for grain or fodder, 0 days for green forage 21-day waiting period 7-day wait for forage, 28 days for grain or fodder 30-day PHI for harvest or grazing 21-day waiting period for harvest Apply in a minimum of 20 gallons of water/acre ground, 3 gallons by air Check label for precautions regarding application of Counter 15G and its interaction with ALS inhibiting herbicides.	
Cruiser 5FS Delta Gold Dimethoate Fastac Force 3G Hero Intrepid Lannate Lorsban 4E Match-Up Mustang MAXX Oberon Onager Poncho Portal Pounce Prevathon Proaxis Radiant Sevin XLR Stallion Warrior II w Zeon Zeal	No grazing restriction 21-day PHI for harvest, 12 days for forage or grazing Apply by aircraft. 14 day PHI for harvest or grazing 30-day PHI for harvest, 60 days for grazing 30-day crop rotation restriction 30-day PHI for grain and stover, 60 days for forage 21-day PHI for harvest 21-day PHI for harvest, 3 days for forage 21-day PHI for harvest, 30-day PHI for harvest or grazing 7-day PHI for harvest 5-day PHI for green forage and silage, 30 days for grain or stover 30-day PHI for harvest or grazing 45-day PHI for harvest or grazing 14-day PHI 30-day PHI for grazing or harvest 14-day PHI for harvest, 1 day for forage, silage, fodder 21-day PHI for harvest or grazing 28-day PHI for harvest, 3 days for fodder or forage 14-day PHI for grazing, 48 days for harvest 30-day PHI for grain, 60 days for forage 21-day PHI for harvest 21-day PHI	

***MOA group numbers in brackets [#] following the insecticide name are used to designate the mode of action of the insecticide according to the classification system developed by the Insecticide Resistance Action Committee, (IRAC). It is intended to help in the selection of insecticides for preventative resistance management. If you make multiple applications for a specific pest during a growing season, simply select a registered insecticide with a different number for each application. To further delay resistance from developing, integrate other control methods into your pest management programs.**

The pesticide information presented in this publication was current with federal and state regulations at the time of revision. **READ and FOLLOW all LABEL directions.**

<i>Pest, Damage and Treatment Threshold</i>	<i>Insecticide Formulation and [Group]* and (active ingredient)</i>	<i>Rate of Product per Acre or 1,000 ft-row</i>	<i>Comments</i>
White grub	Seed Treatments		
Large, "C" shaped grub with a white body and a brown head. Damage: Feed on developing roots, cause slow growth, stunting and stand loss. Threshold: No reliable thresholds are available. Consider using an at-planting treatment for "suppression" if field has a history of grub problems.	Cruiser 5FS [4A] (thiamethoxam)	0.56 to 3.61 fl oz / 80,000 seed	Do not use treated seed for feed, food or oil processing.
	Poncho 600 [4A] (clothianidin)	1.13 fl oz/80,000 seed	Do not use treated seed for feed, food or oil processing.
	Force ST [3]	3 to 4 oz/cwt seed	Do not use Force 3G if Force ST was used. Do not use treated seed for feed, food or oil processing.
	Planting Time		
	Aztec 2.1 G [1B,3] (tebupirromphos, cyfluthrin)	6.7 fl oz/1,000 ft-row	Follow manufactures' guidelines for rates, application methods grazing and crop rotation restrictions. Rotation of insecticides during successive years is suggested.
	Ballista LFC [3] (lambda-cyhalothrin)	0.66 fl oz/1,000 ft-row	
	Counter 15G [1B] (terbufos)	6 to 8 oz/1,000 ft-row	T-band or in-furrow.
	Force 3G [3] (tebupirromphos, cyfluthrin)	4 to 5 oz/1,000 ft-row	T-band or in-furrow.
	Proaxis 0.5 SC [3] (gamma-cyhalothrin)	0.66 fl oz/1,000 ft-row	
Wireworm	Seed Treatments		
Hard-shelled, smooth, cylindrical, yellowish to brown worms. Two- to six-year life cycle. More common in corn planted into a sod or grass pasture. Damage: Feed on seed, seedling. Cause stunting and stand loss. Threshold: No reliable thresholds are available. Treat if field has a history of problems. Wireworms may be more of a problem in no-till or minimum till fields.	Cruiser 5FS [4A] (thiamethoxam)	0.56 to 3.61 fl oz / 80,000 seed	Do not use treated seed for feed, food or oil processing.
	Poncho 600 [4A] (clothianidin)	1.13-2.36 fl oz/ 80,000 seed	Do not use treated seed for feed, food or oil processing.
	Force ST [3] (tebupirromphos + cyfluthrin)	3 to 4 oz/cwt seed	Do not use Force 3G if Force ST was used. Do not use treated seed for feed, food or oil processing.
	Planting Time		
	Aztec 2.1 G [1B,3] (tebupirromphos, cyfluthrin)	6.7 fl oz/1,000 ft-row	Follow manufactures' guidelines for rates, application methods grazing and crop rotation restrictions. Rotation of insecticides during successive years is suggested.
	Ballista LFC [3] (lambda-cyhalothrin)	0.66 fl oz/1,000 ft-row	
	Capture 1.5 G [3]	3.2 to 8 oz/1,000 ft-row	T-band or in-furrow.
	Counter 15G [1B] (terbufos)	6 to 8 oz/1,000 ft-row	T-band or in-furrow.
	Force 3G [3] (tebupirromphos + cyfluthrin)	4 to 5 oz/1,000 ft-row	
	Lorsban 15 G[1B] (chlorpyrifos)	8 oz/1,000 ft-row	
	Proaxis 0.5 SC [3] (gamma-cyhalothrin)	0.66 fl oz/1,000 ft-row	

<i>Pest, Damage and Treatment Threshold</i>	<i>Insecticide Formulation and [Group]* and (active ingredient)</i>	<i>Rate of Product per Acre or 1,000 ft-row</i>	<i>Comments</i>
Corn rootworm (larvae)	Seed Treatments		
Thin, white worm-like larva that lives in soil. Damage is likely to occur in early part of growing season (before June 15). Damage: Feed on roots, causing lodged plants and plants that "gooseneck". Root tissue and brace roots are often chewed back to the base of the stalk. Threshold: Consider a planting-time insecticide, or a seed variety that contains transgenic "root-worm" protection if planting continuous corn.	Rootworm resistant varieties	*Transgenic seed	*Follow company's guidelines for providing refugia, crop rotations and other resistance management strategies.
	Cruiser 5FS [4A] (thiamethoxam)	**5.6 fl oz/80,000 seed	**Do not use treated seed for feed, food or oil processing. See label for mixing and handling instructions. Follow all label restrictions.
	Gaucha 600 [4A]	**6.0 fl oz/80,000 seed	
	Poncho 600 [4A] (clothianidin)	**5.64 fl oz/80,000 seed	
	Force ST [3] (tefluthrin)	3 to 4 oz/cwt seed	Do not use treated seed for feed, food or oil processing. Do not apply Force 3G if Force ST was used.
	Planting Time		
	Aztec 2.1 G [1B,3] (tebupirromphos + cyfluthrin)	6.7 fl oz/1,000 ft-row	Follow manufactures' guidelines for rates, application methods grazing and crop rotation restrictions. Rotation of insecticides during successive years is suggested. Do not make a foliar application if planting time application was made.
	Capture LFR [3] (bifenthrin)	0.39 to 0.98 fl oz/ 1,000 ft-row	
	Counter 15G [1B] (terbufos)	6 to 8 oz/1,000 ft-row	
	Force 3G [3] Force CS [3] (tefluthrin)	4 to 5 oz/1,000 ft-row 0.46 to 0.57 fl oz/ 1,000 ft row	T-band or in-furrow T-band or in-furrow
	Lorsban 15 G [1B] (chlorpyrifos)	2.5 fl oz/1,000 ft-row	T-band or in-furrow
	Proaxis 0.5 CS [3] (gamma-cyhalothrin)	8 oz/1,000 ft-row	
	Thimet 20G [1B] (phorate)	0.24 oz/1,000 ft-row	
	Post Seedling-Emergence Application		
	Counter 15G [1B] (terbufos)	8 oz/1,000 ft-row	Follow label directions for at-cultivation applications. Do not make application if planting time application was made.
	Cobalt [1B,3] (chlorpyrifos+ lambda-cyhalothrin)	38 to 42 fl oz	
	Force 3G [3] (tefluthrin)	4 to 5 oz/1,000 ft-row	
	Lorsban 15 G [1B] (chlorpyrifos)	8 oz/1,000 ft-row	
	Thimet 20G [1B]	4.5 to 6 oz/1,000 ft row	

<i>Pest, Damage and Treatment Threshold</i>	<i>Insecticide Formulation and [Group]* and (active ingredient)</i>	<i>Rate of Product per Acre or 1,000 ft-row</i>	<i>Comments</i>	
<p>Cutworms (black, granulate, sandhill)</p> <p>Striped or solid colored, robust caterpillars that “roll” up when disturbed and prefer to live underground.</p> <p>Damage: Cutworms generally feed at night and live under the soil during the day. Plants will be cut at or slightly above the soil level, causing stand reductions.</p> <p>Threshold: Scout fields at seedling emergence. Treat when worms are less than ½ inch long and skips are noticed.</p>	Seed Treatments			
	Resistant varieties	Transgenic seed	Follow company’s guidelines for providing refugia, crop rotation and other resistance management strategies.	
	Pre-Plant/Planting Time			
	Aztec 2.1 G [1B, 3] (tebuipromphos + cyfluthrin)	6.7 fl oz/1,000 ft-row	Follow manufactures’ guidelines for rates, application methods grazing and crop rotation restrictions.	
	Capture 2EC [3] (bifenthrin)	0.15 to 0.3 fl oz/1,000 ft-row		
	Counter 15G [1B] (terbufos)	6 to 8 oz/1,000 ft-row		
	Force 3G [3] Force CS [3] (tefluthrin)	4 to 5 oz./1,000 ft-row 0.46 to 0.57 fl oz/1,000 ft row	T band or In-furrow T band or In-furrow	
	Lorsban 15 G [1B] (chlorpyrifos)	8 oz/1,000 ft-row	T band or In-furrow	
	Proaxis 0.5 CS [3] (gamma-cyhalothrin)	0.66 oz/1,000 ft-row		
	Pounce 1.5 G [3] (permethrin)	8 oz/1,000 ft-row		
	Post-emergence Sprays			
	Asana XL [3] (esfenvalerate)	5.8 to 9.6 fl oz (0.03 to 0.05 lb ai/A)	21-day waiting period.	
	Baythroid XL [3] (beta-cyfluthrin)	0.8 to 1.6 fl oz (0.007 to 0.013 lb ai/A)	21-day wait for grain or fodder, 0 days for green forage.	
	Besiege[28,3] (chlorantraniliprole + lambda-cyhalothrin)	5.0 to 10 fl oz	21-day waiting period.	
	Brigade 2EC [3] (bifenthrin)	2.1 to 6.4 fl oz (0.033 to 0.1 lb ai/A)	30-day wait for harvest or grazing.	
	Cobalt [1B,3] (chlorpyrifos + gamma-cyhalothrin)	19 to 38 fl oz	21-day wait for harvest.	
	Delta Gold [3] (deltamethrin)	1.0 to 1.5 fl oz (0.012 to 0.018 lb ai/A)	21-day wait for harvest, 12 days for grazing or silage.	
	Fastac [3] (alpha-cypermethrin)	1.3 to 2.8 fl oz (0.008 to 0.018 lb ai/A)	30-day wait for harvest, 60 days for grazing.	
	Hero [3] (zeta-cypermethrin + bifenthrin)	2.6 to 6.1 fl oz	30-day PHI for grain and stover, 60 days for forage.	
	Lorsban 4E [1B] (chlorpyrifos)	1 to 2 pt (0.5 to 1.0 lb ai/A)	21-day wait for grain or harvest.	
Match-Up [1B,3] (chlorpyrifos + bifenthrin)	5.5 to 16.4 fl oz	30-day waiting period for grain and silage, 60 days for grazing.		
Mustang MAXX EC [3] (zeta-cypermethrin)	1.28 to 2.8 fl oz (0.008 to 0.0175 lb ai/A)	21-day waiting period.		
Proaxis 0.5 SC [3] (gamma-cyhalothrin)	1.92 to 3.2 fl oz (0.0075 to 0.0125 lb ai/A)	21-day waiting period for harvest or grazing.		
Pounce 25W [3] (permethrin)	6.4 to 9.6 oz (0.1 to 0.2 lb ai/A)	30-day waiting period.		

CR-7192.6

<i>Pest, Damage and Treatment Threshold</i>	<i>Insecticide Formulation and [Group]* and (active ingredient)</i>	<i>Rate of Product per Acre or 1,000 ft-row</i>	<i>Comments</i>	
<p>Western bean cutworm</p> <p>Larvae are dark brown with faint diamond-shaped markings on their backs. Measures 1.5 inches. Eggs are deposited in masses of 4 to 200 on upper surface of leaves.</p> <p>Damage: Larvae feed on developing tassel or silk. They feed on developing kernels once the ear has formed.</p> <p>Threshold: Treat of 8% or more of the plants have egg masses or small larvae in the tassels and the crop is 95% tasseled.</p>	Seed Treatments			
	Resistant varieties	Transgenic seed	Follow company’s guidelines for providing refugia, crop rotation and other resistance management strategies.	
	Post-emergence Sprays			
	Asana XL [3] (esfenvalerate)	2.9 to 5.8 fl oz (0.015 to 0.03 lb ai/A)	21-day waiting period.	
	Baythroid XL [3] (beta-cyfluthrin)	1.6 to 2.8 fl oz (0.013 to 0.022 lb ai/A)	21-day waiting period for grain or fodder, 0 days for green forage.	
	Besiege[28,3] (chlorantraniliprole + lambda-cyhalothrin)	5.0 to 10 fl oz	21-day wait for harvest.	
	Blackhawk [5] (spinosad)	2.2 to 3.3 fl oz (0.05 to 0.075 lb ai/A)	7-day wait for forage, 28 days for grain or fodder.	
	Brigade 2EC [3] (bifenthrin)	2.6 to 6.4 fl oz (0.033 to 0.1 lb ai/A)	30-day wait for grazing or harvest.	
	Cobalt [1B, 3] (chlorpyrifos + gamma-cyhalothrin)	13 to 26 fl oz	21-day wait for harvest, 14 days for grazing or silage.	
	Coragen [28] (chlorantraniliprole)	3.5 to 7.5 fl oz (0.045 to 0.098 lb ai/A)	14-day waiting period for harvest 1 day for forage, silage, stover.	
	Fastac [3] (alpha-cypermethrin)	1.8 to 3.8 fl oz (0.011 to 0.025 lb ai/A)	30-day wait for harvest, 60 days for grazing.	
	Intrepid 2F [18] (methoxyfenozide)	4 to 16 fl oz (0.06 to 0.25 lb ai/A)	21-day wait for harvest.	
	Karate w Zeon [3] (lambda-cyhalothrin)	0.96 to 1.60 fl oz (0.015 to 0.025 lb ai/A)	21-day wait for harvest or grazing.	
	Lorsban 4E [1B] (chlorpyrifos)	1 to 2 pt (0.5 to 1 lb ai/A)	21-day wait for harvest.	
	Match-Up [1B,3] (chlorpyrifos + bifenthrin)	5.5 to 16.4 fl oz	30-day wait for harvest or grazing.	
	Mustang MAXX EC [3] (zeta-cypermethrin)	1.76 to 4.0 fl oz (0.011 to 0.025 lb ai/A)	7-day waiting period for harvest.	
	Proaxis 0.5 SC [3] (gamma-cyhalothrin)	1.92 to 3.2 fl oz (0.0075 to 0.0125 lb ai/A)	21-day waiting period.	
	Pounce 25 WP (permethrin)	3.2 to 6.4 oz (0.5 to 0.1 lb ai/A)	30-day waiting period.	
	Radiant SC [5] (spinetoram)	3.0 to 6.0 fl oz (0.023 to 0.047 lb ai/A)	28-day wait for harvest, 3 days for forage or fodder.	
	Sevin XLR [1A] (carbaryl)	2 qt (1 lb ai/A)	48-day wait for grain, 14 days for forage or silage.	
Stallion [1B,3] (chlorpyrifos + zeta-cypermethrin)	5.0 to 11.75 fl oz	30-day waiting period for grain, 60 days for forage.		
Warrior II w Zeon [3] (lambda-cyhalothrin)	0.96 to 1.60 fl oz (0.015 to 0.025 lb ai/A)	21-day wait for harvest. Check label for grazing restriction.		

CR-7192.11

<i>Pest, Damage and Treatment Threshold</i>	<i>Insecticide Formulation and [Group]* and (active ingredient)</i>	<i>Rate of Product per Acre or 1,000 ft-row</i>	<i>Comments</i>
Seedcorn maggot, Seed corn beetle (cont'd) Threshold: Replanting is the only recourse if damage has already occurred. Use a planting-time treatment if fields have a history. No-till fields may be more vulnerable to attack.	Capture LFR [3] (bifenthrin)	0.2 to 0.78 fl oz/ 1,000 ft-row	Seed corn beetle only.
	Counter 15G [1B] (terbufos)	6 to 8 oz/1,000 ft-row	
	Lorsban 15G [1B] (chlorpyrifos)	8 to 12 oz/1,000 ft-row	T-band or In-furrow.
	Force 3G [3] (tefluthrin)	4 to 5 oz/1,000 ft-row	T-band or in-furrow.
Southwestern corn borer	Seed Treatments		
Full grown caterpillars are white with prominent dark spots on body. Eggs are laid in masses of 12 to 30. They overlap like egg scales. Eggs are white when first laid, then red bands appear before they hatch.	Resistant varieties	Transgenic seed	Follow company's guidelines for providing refugia, crop rotation and other resistance management strategies.
	Post-emergence Sprays		
Damage: First generation causes "dead heart" in plants. Second generation tunnels throughout stalk. May girdle mature stalks causing lodging.	Baythroid XL [3] (beta-cyfluthrin)	1.6 to 2.8 fl oz (0.013 to 0.022 lb ai/A)	21-day wait for grain or fodder, 0 days for green forage.
	Besiege[28,3] (chlorantraniliprole + lambda-cyhalothrin)	6.0 to 10 fl oz	21-day waiting period.
Threshold: Threshold based on egg masses. Treat if 25% of plants have egg masses or newly hatched larvae. A repeat application may be needed in 7 to 10 days.	Blackhawk [5] (spinosad)	2.2 to 3.3 fl oz (0.05 to 0.075 lb ai/A)	7-day wait for forage, 28 days for grain or fodder.
	Brigade 2EC [3] (bifenthrin)	2.6 to 6.4 fl oz (0.033 to 0.1 lb ai/A)	30-day waiting period for grazing or harvest.
	Cobalt [1B,3] (chlorpyrifos + gamma-cyhalothrin)	19 to 38 fl oz	21-day waiting period for harvest.
	Delta Gold [3] (deltamethrin)	1.5 to 1.9 fl oz (0.018 to 0.022 lb ai/A)	21-day wait for harvest; 12 days for forage or grazing.
	Fastac [3] (alpha-cypermethrin)	2.7 to 3.8 fl oz (0.017 to 0.025 lb ai/A)	30-day wait for grain, 60 days for silage.
	Intrepid 2F [18] (methoxyfenozide)	4 to 16 fl oz (0.06 to 0.25 lb ai/A)	21-day waiting period.
	Karate w Zeon (lambda-cyhalothrin)	1.28 to 1.92 fl oz (0.02 to 0.03 lb ai/A)	35-day waiting period.
	Hero [3] (zeta-cypermethrin + bifenthrin)	4.0 to 10.3 fl oz	30-day PHI for grain and stover, 60 days for forage.
	Lorsban 4E [1B] (chlorpyrifos)	1.5 to 2 pt (0.75 to 1 lb ai/A)	21-day PHI for harvest.
	Match-Up [1B,3] (chlorpyrifos + bifenthrin)	5.5 to 16.4 fl oz	30-day wait for grain or harvest.
	Mustang MAXX EC [3] (zeta-cypermethrin)	2.72 to 4.0 fl oz (0.017 to 0.025 lb ai/A)	7-day waiting period for harvest.
	Prevathon [28] (chlorantraniliprole)	14 to 20 fl oz (0.047 to 0.067 lb ai/A)	14-day waiting period for harvest 1 day for forage, silage, stover.
	Proaxis 0.5 SC [3] (gamma-cyhalothrin)	2.56 to 3.84 fl oz (0.01 to 0.015 lb ai/A)	21-day waiting period.
	Radiant SC [5] (spinetoram)	3.0 to 6.0 fl oz (0.023 to 0.047 lb ai/A)	28-day wait for harvest, 3 days for forage or fodder.
	Stallion [1B,3] (chlorpyrifos + zeta-cypermethrin)	9.25 to 11.75 fl oz	30-day waiting period for grain, 60 days for forage.
	Warrior II w Zeon [3] (lambda-cyhalothrin)	1.28 to 1.92 fl oz (0.02 to 0.03 lb ai/A)	21-day wait for harvest. Check label for grazing restriction.

<i>Pest, Damage and Treatment Threshold</i>	<i>Insecticide Formulation and [Group]* and (active ingredient)</i>	<i>Rate of Product per Acre or 1,000 ft-row</i>	<i>Comments</i>
Cutworms (black, granulate, sandhill) (cont'd)	Stallion [1B,3] (chlorpyrifos + zeta cypermethrin)	3.75 to 11.75 fl oz	30-day waiting period for grain, 60 days for forage.
	Warrior II w Zeon [3] (lambda-cyhalothrin)	0.96 to 1.60 fl oz (0.015 to 0.025 lb ai/A)	21-day wait for harvest. Check label for grazing restriction.
Fall armyworm	Seed Treatments		
Large, striped, non-bristled worm up to 1.5 inches. Has a light colored, inverted "Y" on head. June-August	Resistant varieties	Transgenic seed	Follow company's guidelines for providing refugia, crop rotation and other resistance management strategies.
	Post-emergence Sprays		
Damage: Larvae cut holes in leaves at whorl stage, heaviest damage occurs on late corn when caterpillars tunnel into ear or ear shank.	Baythroid XL [3] (beta-cyfluthrin)	2.8 fl oz (0.022 lb ai/A)	21-day wait for grain or fodder, 0 days for green forage.
	Besiege[28,3] (chlorantraniliprole + lambda-cyhalothrin)	6.0 to 10 fl oz	21-day waiting period.
Threshold: Treat if 75% of plants are infested during whorl stage.	Blackhawk [5] (spinosad)	1.67 to 3.3 fl oz (0.038 to 0.075 lb ai/A)	7-day wait for forage, 28 days for grain or fodder.
	Brigade 2EC [3] (bifenthrin)	2.1 to 6.4 fl oz (0.033 to 0.1 lb ai/A)	30-day wait for grazing or harvest.
	Cobalt [1B, 3] (chlorpyrifos + gamma-cyhalothrin)	13 to 26 fl oz	21-day waiting period for harvest, 14 days for grazing or silage.
	Coragen [28] (chlorantraniliprole)	3.5 to 7.5 fl oz (0.045 to 0.098 lb ai/A)	14-day waiting period for harvest 1 day for forage, silage, stover.
	Delta Gold [3] (deltamethrin)	1.5 to 1.9 fl oz (0.018-0.022 lb ai/A)	21-day wait for harvest, 12 days for grazing or silage.
	Fastac [3] (alpha-cypermethrin)	3.2 to 3.8 fl oz (0.020 to 0.025 lb ai/A)	30-day wait for harvest, 60 days for grazing.
	Hero [3] (zeta-cypermethrin + bifenthrin)	4.0 to 10.3 fl oz	30-day PHI for grain and stover, 60 days for forage.
	Lannate LV [1A] (methomyl)	0.75 to 1.5 pt 0.225 to 0.45 lb	21-day PHI for harvest, 3 days for forage.
	Lorsban 4E [1B] (chlorpyrifos)	1 to 2 pt (0.5 to 1.0 lb ai/A)	21-day waiting period.
	Match-Up [1b,3] (chlorpyrifos + bifenthrin)		30-day wait for grain.
	Mustang MAXX EC [3] (zeta-cypermethrin)	3.2 to 4.0 fl oz (0.02 to 0.025 lb ai/A)	7-day waiting period for harvest.
	Pounce 25 WP [3] (permethrin)	6.4 to 9.6 fl oz (0.1 to 0.15 lb ai/A)	30-day waiting period.
	Proaxis 0.5 SC [3] (gamma-cyhalothrin)	2.56 to 3.84 fl oz (0.01 to 0.015 lb ai/A)	21-day waiting period.
	Stallion[1B, 3] (chlorpyrifos + zeta-cypermethrin)	9.25 to 11.75 fl oz	30-day waiting period for grain, 60 days for forage.
	Warrior II w Zeon [3] (lambda-cyhalothrin)	1.28 to 1.92 fl oz (0.02 to 0.03 lb ai/A)	21-day waiting period. Check label for grazing restrictions.

<i>Pest, Damage and Treatment Threshold</i>	<i>Insecticide Formulation and [Group]* and (active ingredient)</i>	<i>Rate of Product per Acre or 1,000 ft-row</i>	<i>Comments</i>
Flea beetles	Asana XL [3] (esfenvalerate)	5.8 to 9.6 fl oz (0.03 to 0.05 lb ai/A)	21-day waiting period.
Shiny, black beetle about 1/16 inch that jumps when disturbed.	Baythroid XL [3] (beta-cyfluthrin)	0.8 to 1.6 fl oz (0.007 to 0.013 lb ai/A)	21-day wait for grain or fodder, 0 days for green forage.
Damage: Early spring-summer. Plant tissue is scraped from leaf, giving it a drought-stressed appearance. Can cause delayed development in cool growing conditions.	Besiege[28,3] (chlorantraniliprole + lambda-cyhalothrin)	6.0 to 10 fl oz	21-day waiting period.
Threshold: Apply to small plants when beetles first appear and some plants are being killed.	Brigade 2EC [3] (bifenthrin)	2.1 to 6.4 fl oz (0.033 to 0.1 lb ai/A)	30-day wait for grazing or harvest.
	Cobalt [1B, 3] (chlorpyrifos + gamma-cyhalothrin)	13 to 26 fl oz	21-day waiting period for harvest.
	Delta Gold [3] (deltamethrin)	1.0 to 1.5 fl oz (0.012 to 0.018 lb ai/A)	21-day wait for harvest, 12 days for grazing or silage.
	Fastac [3] (alpha-cypermethrin)	2.7 to 3.8 fl oz (0.017 to 0.025 lb ai/A)	30-day wait for grain, 60 days for silage.
	Hero [3] (zeta-cypermethrin + bifenthrin)	2.6 to 6.1 fl oz	30-day wait for grain and silage, 60 days for grazing.
	Lannate LV [1A] (methomyl)	0.75 to 1.5 pt 0.225 to 0.45 lb	21-day PHI for harvest, 3 days for forage.
	Lorsban 4E [1B] (chlorpyrifos)	1 to 2 pt (0.5 to 1.0 lb ai/A)	21-day waiting period.
	Match-Up [1B,3] (chlorpyrifos + bifenthrin)	5.5 to 16.4 fl oz	30-day wait for grain or harvest.
	Mustang MAXX EC [3] (zeta-cypermethrin)	2.72 to 4.0 fl oz (0.017 to 0.025 lb ai/A)	30-day wait for grain and silage, 60 days for grazing.
	Proaxis 0.5 SC [3] (gamma-cyhalothrin)	2.56 to 3.84 fl oz (0.01 to 0.015 lb ai/A)	21-day waiting period.
	Pounce 25WP [3] (permethrin)	6.4 to 9.6 oz (0.1 to 0.2 lb ai/A)	30-day waiting period.
	Sevin XLR [1A] (carbaryl)	1 to 2 qt (0.5 to 1 lb ai/A)	48-day wait for grain, 14 days for forage or silage.
	Stallion[1B,3] (chlorpyrifos + zeta-cypermethrin)	9.25 to 11.75 fl oz	30-day waiting period for grain, 60 days for forage.
	Warrior II w Zeon [3] (lambda-cyhalothrin)	1.28 to 1.92 fl oz (0.02 to 0.03 lb ai/A)	21-days wait for harvest. Check label for grazing restriction.
Grasshopper	Asana XL [3] (esfenvalerate)	2.9 to 5.8 fl oz (0.015 to 0.03 lb ai/A)	21-day waiting period.
1-2 inches, outer wings leathery, inner wings clear or colored. Enlarged hind legs designed for jumping.	Baythroid XL [3] (beta-cyfluthrin)	2.1 to 2.8 fl oz (0.017 to 0.022 lb ai/A)	21-day wait for grain or fodder, 0 day for green forage.
Damage: Chew leaves, leaving ragged edges, or completely chewing leaf blade. Damage emerging seed heads, causing yield loss.	Brigade 2EC [3] (bifenthrin)	2.1 to 6.4 fl oz (0.033 to 0.1 lb ai/A)	30-day wait for grazing or harvest.
Threshold: Consider treating if numbers reach 8 to 14 in the field, or 20 to 40 in field margins.	Cobalt [1B, 3] (chlorpyrifos + gamma-cyhalothrin)	7 to 13 fl oz	21-day waiting period for harvest, 14 days for grazing or silage.
	Coragen [28] (chlorantraniliprole)	2.0 to 5.0 fl oz (0.026 to 0.065 lb ai/A)	14-day waiting period for harvest 1 day for forage, silage, stover.
	Delta Gold [3] (deltamethrin)	1 to 1.5 fl oz (0.012-0.018 lb ai/A)	21-day wait for harvest; 12 days for grazing or forage.

CR-7192.8

<i>Pest, Damage and Treatment Threshold</i>	<i>Insecticide Formulation and [Group]* and (active ingredient)</i>	<i>Rate of Product per Acre or 1,000 ft-row</i>	<i>Comments</i>
Grasshopper (cont'd) See F-7196, Grasshopper Management in Rangeland, Pastures and Crops for more information.	Fastac [3] (alpha-cypermethrin)	2.7 to 3.8 fl oz (0.017 to 0.025 lb ai/A)	30-day wait for harvest, 60 days for grazing.
	Hero [3] (zeta-cypermethrin + bifenthrin)	2.6 to 6.1 fl oz	30-day PHI for grain and stover, 60 days for forage.
	Karate w Zeon [3] (lambda-cyhalothrin)	1.28 to 1.92 fl oz (0.02 to 0.03 lb ai/A)	21-day waiting period.
	Lorsban 4E [1B] (chlorpyrifos)	1 to 2 pt (0.5 to 1 lb ai/A)	21-day waiting period.
	Match-Up [1B,3] (chlorpyrifos + bifenthrin)	5.5 to 16.4 fl oz	30-day wait for harvest or grazing.
	Mustang MAXX EC [3] (zeta-cypermethrin)	2.72 to 4.0 fl oz (0.017 to 0.025 lb ai/A)	7-day waiting period for harvest.
	Proaxis 0.5 SC [3] (gamma-cyhalothrin)	1.92 to 3.2 fl oz (0.0075 to 0.0125 lb ai/A)	21-day waiting period.
	Warrior II w Zeon [3] (lambda-cyhalothrin)	1.28 to 1.92 fl oz (0.02 to 0.03 lb ai/A)	21-day wait for harvest. Check label for grazing restriction.
Mites	Capture 2EC [3] (bifenthrin)	5.12 to 6.4 fl oz (0.08 to 0.1 lb ai/A)	30-day waiting period.
Small, less than 1/100 inch. Cause brown stippling of leaves. Banks grass and two-spotted spidermites are most common pests.	Comite II [20] (propargite)	36 to 54 fl oz/Acre	30-day waiting period. Apply when mite colonies first form, when leaves are dry.
	Dimethoate 4E [1B]	0.66 to 1 pt	14-day waiting period.
Damage: Causes stippling of leaves, severe infestations can kill leaves. Infestations generally start at lower leaves and move upward.	Hero [3] (zeta-cypermethrin + bifenthrin)	10.3 fl oz	30-day wait for grain, 60 days for silage.
Threshold: Treat when there is visible damage on the lower third of the plant and small colonies are visible on the middle third of the plant, and the corn has not yet reached the hard dough stage.	Oberon 2 SC [23] (spiromesifen)	2.85 to 8.0 fl oz	5-day PHI for green forage and silage, 30 days for grain or stover.
	Onager [10A] (hexythiazox)	10 to 24 fl oz	30-day waiting period.
	Portal [21A] (fenpyroximate)	2 pints 0.1 lb ai/A)	14-day waiting period.
	Zeal WDG [10B] (etoxazole)	1.0 to 3.0 oz (0.045 to 0.135 lb ai/A)	* for seed production only, 21-day waiting period.
			NOTE: Treatments at hard-dough stage or later are not cost effective. When heavy infestations occur, erratic control will usually be the rule. Thorough cover-age is important, higher volumes (2 to 3 gallons or more per acre) when applied by aircraft increase the effectiveness of the spray.
Seedcorn maggot, Seed corn beetle	Seed Treatments		Follow manufactures' guidelines for rates, application methods grazing and crop rotation restrictions. Rotation of insecticides during successive years is suggested.
Maggots are yellowish-white, tapered larvae about 1/4 inch. Beetles are about 3/8 inch, with two black stripes on brown wing covers.	Cruiser 5FS [4A] (thiamethoxam)	0.56 to 3.61 fl oz / 80,000 seed	
	Poncho 600 [4A] (clothianidin)	1.13 to 2.26 fl oz/ 80,000 seed	
	Force ST [3] (tefluthrin)	3 to 4 oz/cwt seed	
Damage: Damage occurs in spring, especially if soils are cool and moist and seeds are not germinating rapidly. Damage is notices as skips in plant stands. Seed will be hollowed out.	Planting Time		
	Aztec 2.1 G [1B,3] (tebupirromphos, cyfluthrin)	6.7 fl oz/1,000 ft-row	

CR-7192.9