



Background Metal Concentrations in Oklahoma Soils

The Oklahoma Cooperative Extension Service Bringing the University to You!

The Cooperative Extension Service is the largest, most successful informal educational organization in the world. It is a nationwide system funded and guided by a partnership of federal, state, and local governments that delivers information to help people help themselves through the land-grant university system.

Extension carries out programs in the broad categories of agriculture, natural resources and environment; family and consumer sciences; 4-H and other youth; and community resource development. Extension staff members live and work among the people they serve to help stimulate and educate Americans to plan ahead and cope with their problems.

Some characteristics of the Cooperative Extension system are:

- The federal, state, and local governments cooperatively share in its financial support and program direction.
- It is administered by the land-grant university as designated by the state legislature through an Extension director.
- Extension programs are nonpolitical, objective, and research-based information.
- It provides practical, problem-oriented education

for people of all ages. It is designated to take the knowledge of the university to those persons who do not or cannot participate in the formal classroom instruction of the university.

- It utilizes research from university, government, and other sources to help people make their own decisions.
- More than a million volunteers help multiply the impact of the Extension professional staff.
- It dispenses no funds to the public.
- It is not a regulatory agency, but it does inform people of regulations and of their options in meeting them.
- Local programs are developed and carried out in full recognition of national problems and goals.
- The Extension staff educates people through personal contacts, meetings, demonstrations, and the mass media.
- Extension has the built-in flexibility to adjust its programs and subject matter to meet new needs. Activities shift from year to year as citizen groups and Extension workers close to the problems advise changes.

Hailin Zhang
Regents Professor and Director
Soil, Water, and Forage Analytical Lab

Trace element concentrations vary naturally in native soils primarily due to differences in mineral composition and the chemical characteristics of the parent material (Adriano, 2001). Anthropogenic sources such as the input of fertilizers, pesticides and other soil amendments such as biosolids, animal manures and drilling mud may increase trace element levels in soils. For example, lead arsenate was used on cotton fields and orchards as a pesticide in the 1940s. Other activities that have affected metal concentrations in soils of Oklahoma include mining and smelting, emissions from different industries and the deposition of vehicle exhausts.

Metals may become an environmental problem when present at elevated concentrations due to their persistence, tendency to bioaccumulate, mobility and potential toxicity in the environment. Background metal concentrations of soils are often used as criteria for assessing environmental quality. Background levels must be determined to establish accurate and realistic guideline levels. Accurate background concentrations of metal and other trace elements allow regulators and consultants to make more informed decisions on the

Oklahoma Cooperative Extension Fact Sheets are also available on our website at: <http://osufacts.okstate.edu>

remediation of different soils with varying properties. Twenty-eight Oklahoma benchmark soils were collected from various locations representing major land resource areas across the state to establish the background levels of metals and trace elements in the state. The locations of the sampling sites are shown in Figure 1. The soils had no known history of applications of fertilizers, biosolids or manure within three years of collection. The soils were digested according to USEPA Method 3050B and analyzed for eight metals, one metalloid (arsenic) and one nonmetal (selenium) using inductively coupled plasma-atomic emission spectroscopy (ICP-AES). The concentrations of those elements (referred as metals thereafter) in the 28 benchmark soils are presented in Table 1. The concentrations of each element vary widely among different soils. Therefore, the median value presented at the bottom of the table would serve as a better background level for the state.




Figure 1. The delineation of major land resource areas in Oklahoma and the locations of the 28 benchmark soils used to establish the background metal concentrations (Scott, 1994; Zhang et al., 2005, Richards et al., 2012). The benchmark soils represent the diversity of soils in Oklahoma as well as the Major Land Resource Areas.

Oklahoma State University, in compliance with Title VI and VII of the Civil Rights Act of 1964, Executive Order 11246 as amended, and Title IX of the Education Amendments of 1972 (Higher Education Act), the Americans with Disabilities Act of 1990, and other federal and state laws and regulations, does not discriminate on the basis of race, color, national origin, genetic information, sex, age, sexual orientation, gender identity, religion, disability, or status as a veteran, in any of its policies, practices or procedures. This provision includes, but is not limited to admissions, employment, financial aid, and educational services. The Director of Equal Opportunity, 408 Whitehurst, OSU, Stillwater, OK 74078-1035; Phone 405-744-5371; email: eeo@okstate.edu has been designated to handle inquiries regarding non-discrimination policies: Director of Equal Opportunity. Any person (student, faculty, or staff) who believes that discriminatory practices have been engaged in based on gender may discuss his or her concerns and file informal or formal complaints of possible violations of Title IX with OSU's Title IX Coordinator 405-744-9154.

Issued in furtherance of Cooperative Extension work, acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture, Director of Oklahoma Cooperative Extension Service, Oklahoma State University, Stillwater, Oklahoma. This publication is printed and issued by Oklahoma State University as authorized by the Vice President for Agricultural Programs and has been prepared and distributed at a cost of 20 cents per copy. Revised 0518. GH.

Table 1. Trace elements concentrations (parts per million or ppm, mg/kg) in the 28 benchmark soils of Oklahoma (Richards et al., 2012).

Soil	As	Cd	Cr	Cu	Mn	Mo	Ni	Pb	Se	Zn
Bernow	0.75	0.13 [†]	4.83	2.13	27.4	0.31	2.41	5.45	<1.25	15.3
Burleson	4.93	0.36	53.7	14.3	580	1.25	21.6	19.1	<1.25	67.6
Carnasaw	12.8	0.66	51.0	8.43	1690	1.80	25.1	29.0	<1.25	63.2
Clarksville	2.95	0.30	16.7	11.9	2165	0.69	15.3	19.3	<1.25	59.1
Cobb	2.30	0.13 [†]	10.7	5.78	106	0.26	7.46	6.04	<1.25	26.4
Dalhart	2.01	0.13 [†]	7.14	3.78	98	0.26	5.46	4.23	<1.25	26.5
Darnell	1.13	0.13 [†]	6.95	3.39	236	0.22	4.91	6.46	<1.25	24.6
Dennis	10.3	0.34	32.8	10.7	515	0.83	19.3	12.9	<1.25	53.0
Dougherty	1.08	0.13 [†]	4.63	2.76	133	0.38	2.85	7.81	<1.25	40.8
Durant	4.72	0.13 [†]	25.7	10.7	508	0.38	10.9	17.1	<1.25	45.2
Easpur	5.6	0.28	22.6	10.0	388	0.45	15.2	19.9	<1.25	57.3
Grant	3.19	0.25	20.1	9.36	383	0.46	14.3	11.9	<1.25	47.2
Kirkland	4.84	0.29	27.2	14.5	523	0.53	17.5	16.4	<1.25	61.2
Lebron	10.5	0.75	51.0	31.9	658	1.14	37.2	22.6	<1.25	106
Mansic	4.20	0.40	22.4	16.0	332	0.50	15.5	12.3	<1.25	61.4
Osage	9.16	0.80	54.1	31.2	725	0.84	38.0	27.6	<1.25	142
Parsons	24.4	0.73	57.0	20.1	853	1.29	29.4	28.8	<1.25	94.2
Pond Creek	2.93	0.25	19.6	11.5	409	0.41	13.9	12.0	<1.25	50.6
Pratt	1.45	0.13 [†]	4.30	1.94	52.6	0.18	3.00	2.60	<1.25	20.9
Renfrow	3.73	0.13 [†]	19.9	10.9	292	0.41	12.3	13.7	<1.25	41.7
Richfield	6.29	0.39	29.4	19.4	469	0.74	20.1	15.1	<1.25	79.5
St. Paul	4.18	0.13 [†]	19.9	12.1	314	0.39	14.2	10.2	<1.25	47.8
Sallisaw	3.13	0.13 [†]	19.7	11.3	1475	1.13	12.3	14.3	<1.25	51.7
Stiegler	3.74	0.13 [†]	19.4	32.7	1601	0.53	10.7	20.2	<1.25	56.7
Summit	33.6	0.76	69.7	15.6	2119	1.00	57.3	31.7	<1.25	72.1
Tillman	4.34	0.13 [†]	26.4	13.6	506	0.50	19.7	13.3	<1.25	56.7
Woodward	3.61	0.13 [†]	19.9	10.4	293	0.36	16.2	7.49	<1.25	41.5
Zaneis	3.03	0.13 [†]	16.8	6.43	166	0.43	9.91	9.41	<1.25	35.0
Mean	6.25	0.30	26.2	12.6	629	0.63	16.9	14.9	<1.25	55.2
Median	3.96	0.19	20.0	11.1	439	0.48	14.8	13.5	<1.25	52.4
Minimum	0.75	0.13	4.30	1.94	27.4	0.18	2.41	2.60	<1.25	15.3
Maximum	33.6	0.80	69.7	32.7	2165	1.80	57.3	31.7	<1.25	142

[†] The sample measured below the detection limit (BDL) of 0.25 mg/kg, but for statistical purposes, the BDL samples were replaced with one-half the detection limit. As (arsenic), Cd (cadmium), Cr (chromium), Cu (copper), Mn (manganese), Mo (molybdenum), nickel (Ni), lead (Pb), selenium (Se), and zinc (Zn).

Table 2 compares concentrations of background metals with the current USEPA Region 6 Screening Levels for these metals. The screening levels (SLs) were developed using risk assessment guidance from the EPA Superfund program (RAGS). The SLs are risk-based concentrations derived from standardized equations that combine exposure information and toxicity data and are intended to be protective of human health. The SLs are generic (i.e., they are calculated without site-specific information) and may be re-calculated using site-specific data. The Oklahoma Department of Environmental Quality (ODEQ) uses EPA Region 6 SLs to determine if a particular site requires clean-up. Usually, if metal concentrations are less than the SLs, no further action or study is required at a site (ODEQ, 2013). The industrial SLs apply to a situation which is not residential (i.e. low chance of regular human exposure) and use a different set of assumptions for the exposure information and calculations. The industrial SLs are often useful to consultants involved in environmental work such as site assessments, tank removals and remediation. Overall,

the industrial SLs are not as conservative as the residential SLs. Generally, ODEQ uses the most conservative level for screening purposes. Certain chemicals, including metals, may produce either carcinogenic or noncarcinogenic toxicological effects or both. Therefore, both carcinogenic and noncarcinogenic end points are evaluated during risk assessment and carcinogenic screening levels have been determined for some metals.

The EPA Region 6 Screening Levels were developed jointly with DOE's Oak Ridge National Laboratory (ORNL) under an Interagency Agreement as an update of the EPA Region 3 RBC Table, Region 6 HHMSSL Table and the Region 9 PRG Table.

Both As and Cd presented in Table 2 are known human carcinogens and have established carcinogenic SLs. The carcinogenic USEPA SLs for As residential and industrial exposures are 0.67 mg/kg and 3.0 mg/kg, respectively (using a target cancer risk of 1E-06). The residential carcinogenic SL for Cd is 2,100 mg/kg while the industrial carcinogenic

Table 2. Comparison of background median metal concentrations and ranges of Oklahoma benchmark soils with USEPA Region 6 Screening Level (SL) for residential and industrial exposures.

Trace Element	Median Oklahoma Soil Levels (mg /kg)	Noncarcinogenic Residential SL [†] (mg/kg)	Noncarcinogenic Industrial SL (mg/kg)
Arsenic (As)	3.96 (0.75-33.6)	34	380
Cadmium (Cd)	0.19 (0.13-0.80)	70	980
Chromium (Cr)	20.0 (4.30-69.7)	Not listed	Not listed
Copper (Cu)	11.1 (1.94-32.7)	3,100	47,000
Manganese (Mn)	439 (27.4-2165)	1,800	26,000
Molybdenum (Mo)	0.48 (0.18-1.80)	390	5,800
Nickel (Ni)	14.8 (2.41-57.3)	1,500	22,000
Lead (Pb)	13.5 (2.60-31.7)	400	800
Selenium (Se)	<1.25	390	5,800
Zinc (Zn)	52.4 (15.3-142)	23,000	350,000

[†] Obtained from USEPA Region 6 Regional Screening Levels (Hazard Index = 1) (http://www.epa.gov/reg3hwmd/risk/human/rb-concentration_table/Generic_Tables/index.htm).

SL is 9,300 mg/kg. Notice the background levels of the trace elements listed in Table 2 are considerably less than the residential SL with the exception of As. However, the mean background level of arsenic found in Oklahoma soils is approximately six times greater than the residential SL of 0.67 mg/kg established by the USEPA. The relatively high level of As in Oklahoma soils is probably due to naturally occurring high levels of As in geologic parent materials. It appears the regional SLs established for As are too low for Oklahoma soils and that background levels may be a useful alternative for baseline screenings and other risk assessments. Additionally, the use of background levels in these situations may eliminate the burden of unneeded soil remediation. Currently, the ODEQ recommends that uncontaminated samples be obtained in the area of the site under study and analyzed to determine background level of As that may be used as the screening level (ODEQ, personal communication).

References

Adriano, D.C. 2001. Trace elements in terrestrial environments: Biogeochemistry, bioavailability, and risks of metals. 2nd ed. Springer-Verlag, New York.

Oklahoma Department of Environmental Quality (ODEQ). 2013. Risk-Based Decision Making for Site Cleanup. <http://www.deq.state.ok.us/factsheets/land/SiteCleanup.pdf> (accessed on August 1, 2014)

Richards, J.R., J.L. Schroder, H. Zhang, N.T. Basta, Y. Wang, and M.E. Payton. 2012. Trace elements in benchmark soils of Oklahoma. *Agron. J.* 76:2031–2040.

Scott, T.D. 1994. Distribution, speciation, and bioavailability of heavy metals in soils of Oklahoma. M.S. thesis. Oklahoma State Univ., Stillwater.

USEPA. 1996. Method 3050b: Acid digestion of sediments, sludges, and soils. In: Test methods for evaluating solid waste: Physical/chemical methods. SW-846. USEPA, Washington, DC. Available at <http://www.epa.gov/wastes/hazard/test-methods/sw846/pdfs/3050b.pdf> (accessed on August 1, 2014)

USEPA. 2014. Region 6 Regional Screening Levels (Formerly HHMSSL - Human Health Medium-Specific Screening Levels). http://www.epa.gov/reg3hwmd/risk/human/rb-concentration_table/Generic_Tables/index.htm (accessed on August 1, 2014)

Zhang, H., J. Schroder, J.K. Fuhrman, N.T. Basta, D.E. Storm, and M.E. Payton. 2005. Path and multiple regression analyses of phosphorus sorption capacity. *Soil Sci. Soc. Am. J.* 69:96–106.